

UNITARIZABILITY, MAUREY-NIKISHIN FACTORIZATION, AND POLISH GROUPS OF FINITE TYPE

HIROSHI ANDO, YASUMICHI MATSUZAWA, ANDREAS THOM, AND ASGER TÖRNQUIST

ABSTRACT. Let Γ be a countable discrete group, and let $\pi: \Gamma \rightarrow \text{GL}(H)$ be a representation of Γ by invertible operators on a separable Hilbert space H . We show that the semidirect product group $G = H \rtimes_{\pi} \Gamma$ is SIN (G admits a two-sided invariant metric compatible with its topology) and unitarily representable (G embeds into the unitary group $\mathcal{U}(\ell^2(\mathbb{N}))$), if and only if π is uniformly bounded, and that π is unitarizable if and only if G is of finite type: that is, G embeds into the unitary group of a II_1 -factor. Consequently, we show that a unitarily representable Polish SIN groups need not be of finite type, answering a question of Sorin Popa. The key point in our argument is an equivariant version of the Maurey–Nikishin factorization theorem for continuous maps from a Hilbert space to the space $L^0(X, m)$ of all measurable maps on a probability space.

1. INTRODUCTION

In this paper, we continue our study [AM12-2] of Popa’s class $\mathcal{U}_{\text{fin}}$ of finite type Polish groups. Here, a Polish group G is said to be of *finite type* if it is isomorphic as a topological group to a closed subgroup of the unitary group $\mathcal{U}(M)$ of some finite von Neumann algebra M with separable predual, where $\mathcal{U}(M)$ is endowed with the strong operator topology. The class $\mathcal{U}_{\text{fin}}$ was introduced in [Po07], where Popa proved the celebrated Cocycle Superrigidity Theorem. This theorem asserts if Γ is a w -rigid group and $\alpha: \Gamma \curvearrowright (X, \mu)$ is a probability measure preserving (p.m.p) action of Γ on a standard probability space (X, μ) with certain properties, then any 1-cocycle $w: X \times \Gamma \rightarrow G$ for α with values in a Polish group $G \in \mathcal{U}_{\text{fin}}$ is cohomologous to a group homomorphism $\Gamma \rightarrow G$, see [Po07] for details. In spite of the importance of Popa’s Theorem and its wide range of applications, not much has been known about the properties of the class $\mathcal{U}_{\text{fin}}$. There are two necessary conditions for a Polish group to be in $\mathcal{U}_{\text{fin}}$. Namely, if $G \hookrightarrow \mathcal{U}(M)$ for some finite von Neumann algebra (M, τ) with separable predual, then clearly $G \hookrightarrow \mathcal{U}(\ell^2(\mathbb{N}))$. G is said to be *unitarily representable* if this condition is satisfied. Moreover, the restriction of the 2-norm metric d on $\mathcal{U}(M)$ given by

$$d(u, v) := \|u - v\|_2, \quad \|x\|_2 := \tau(x^*x)^{\frac{1}{2}} \quad (u, v \in \mathcal{U}(M))$$

defines a two-sided invariant metric on G compatible with the topology. It is well-known that a metrizable topological group admits a two-sided invariant metric compatible with the topology if and only if the group is SIN (see Definition 2.3). In [Po07, §6.5], Popa posed a question whether the above two conditions are sufficient:

Question (Popa). Let G be a unitarily representable SIN Polish group. Is G necessarily of finite type?

Key words and phrases. uniformly bounded representation, unitarizability, Maurey–Nikishin factorization, Polish groups of finite type.

Motivated by the study [AM12-1] of the Lie structure of the class $\mathcal{U}_{\text{fin}}$, two of us started a research on the above Popa's $\mathcal{U}_{\text{fin}}$ -problem [AM12-2], and obtained some positive answers under additional conditions on G . Among other things, we showed that the question has a positive answer if G is amenable. For example, the Hilbert-Schmidt group $\mathcal{U}(H)_2$ is a finite type Polish group. However, the general case was left wide open.

Note that there is a similarity between the $\mathcal{U}_{\text{fin}}$ -problem and the problem of finding a trace on a finite von Neumann algebra, which was established by Murray and von Neumann in their classical papers on the subject. Yeadon [Ye71] found a simpler proof of Murray's and von Neumann's theorem as follows: Let M be a finite von Neumann algebra. We assume for simplicity that M has separable predual, and let φ be a faithful normal state on M . We let the unitary group $\mathcal{U}(M)$ act on the predual M_* by conjugation:

$$(1) \quad \alpha_u(\psi) := u\psi u^* = \psi(u^* \cdot u), \quad \psi \in M_*, u \in \mathcal{U}(M).$$

Note that α is an affine isometric action on the Banach space M_* . By using the finiteness assumption, Yeadon showed that the orbit $\mathcal{O}_\varphi := \{\alpha_u(\varphi); u \in \mathcal{U}(M)\}$ is relatively weakly compact. Thus its weak convex closure $\overline{\text{co}}(\mathcal{O}_\varphi) \subset M_*$ is a weakly compact convex subset of M_* on which $\mathcal{U}(M)$ acts by affine isometries. By the Ryll-Nardzewski Theorem, this implies that there exists an α -fixed point, which is easily seen to be a normal faithful tracial state τ .

In the first named author's PhD thesis, we adapted the Yeadon argument up to some point. This part of the thesis is an unpublished joint work with the second named author. Let G be a unitarily representable Polish SIN group. It is well-known (see e.g. [Gao05]) that G is unitarily representable if and only if G admits a continuous positive definite function φ which generates the topology. An analogous characterization for $\mathcal{U}_{\text{fin}}$ is obtained in [AM12-2]: Let $\mathcal{P}(G)$ be the set of all normalized continuous positive definite functions on G . Then G is of finite type if and only if there exists a character (i.e., a function $\varphi \in \mathcal{P}(G)$ which is conjugation-invariant) which generates a neighborhood basis of the identity on G (see Theorem 2.4 below for details). It is clear that $\mathcal{P}(G)$ is a convex subset of the Banach space $C_b(G)$ of all continuous bounded functions on G equipped with the sup norm. Define an affine isometric action $\beta: G \curvearrowright C_b(G)$ by:

$$\beta_g(\psi) := \psi(g^{-1} \cdot g), \quad \psi \in C_b(G), g \in G.$$

If the orbit $\mathcal{O}_\varphi = \{\beta_g(\varphi); g \in G\}$ is relatively weakly compact, then by the Ryll-Nardzewski Theorem, there exists a fixed point ψ in the closed convex hull $\overline{\text{co}}(\mathcal{O}_\varphi)$. Once we get a fixed point $\psi \in \overline{\text{co}}(\mathcal{O}_\varphi)$, one can show, using Theorem 2.4 below, that G is of finite type. Unfortunately, we do not know if $\mathcal{O}_\varphi$ is relatively weakly compact. What we could show is the following result.

Theorem 1.1 (Ando–Matsuzawa, unpublished). *Let G be a unitarily representable Polish SIN group, and let $\varphi \in \mathcal{P}(G)$ generates the topology of G . Then the following statements hold:*

- (i) $\mathcal{O}_\varphi$ is relatively compact in the compact open topology.
- (ii) If $\mathcal{O}_\varphi$ satisfies one of the following conditions, then G is of finite type.
 - (iia) The closure of $\mathcal{O}_\varphi$ with respect to the topology of pointwise convergence is norm-separable.
 - (iib) $\mathcal{O}_\varphi$ is relatively weakly compact.

From the above result, two of us considered that this was an indication that the $\mathcal{U}_{\text{fin}}$ -problem may have a positive answer in general. However, it was not clear how to verify the

relationship between the compact open topology and the weak topology in $C_b(G)$ or to verify either (ia) or (ib).

In this paper, we show that indeed the orbit $\mathcal{O}_\varphi \subset C_b(G)$ is not always relatively weakly compact. That is, we construct a family of unitarily representable SIN Polish groups which fail to be in the class $\mathcal{U}_{\text{fin}}$. This answers the above question of Popa. The construction is based on the existence of uniformly bounded non-unitarizable representations of countable discrete groups:

Theorem 1.2 (Main Theorem). *Let Γ be a countable discrete group, and let $\pi: \Gamma \rightarrow \text{GL}(H)$ be a group homomorphism from Γ to the group of bounded invertible operators on H . Let $G = H \rtimes_\pi \Gamma$ be the semidirect product.*

- (a) *G is a unitarily representable Polish group. Moreover, the following two conditions are equivalent.*
 - (i) *G is a SIN group.*
 - (ii) *π is uniformly bounded. That is, $\|\pi\| := \sup_{s \in \Gamma} \|\pi(s)\| < \infty$ holds.*
- (b) *Assume that π is uniformly bounded. Then the following three conditions are equivalent.*
 - (i) *G is of finite type.*
 - (ii) *There exists a character $f: H \rightarrow \mathbb{C}$ which generates a neighborhood basis of $0 \in H$, such that $f(\pi(s)\xi) = f(\xi)$ for all $\xi \in H$ and $s \in \Gamma$.*
 - (iii) *π is unitarizable. That is, there exists $V \in \text{GL}(H)$ such that $V^{-1}\pi(s)V$ is unitary for all $s \in \Gamma$.*

A group Γ is said to be *unitarizable* if all of its uniformly bounded representations are unitarizable. The problem of unitarizability has a long history. Sz.-Nagy [SzN47] showed that the group $\mathbb{Z}$ of integers is unitarizable. Later, this was generalized to amenable groups independently by Dixmier [Di50], Nakamura–Takeda [NT51] and Day [Da50]. This led Dixmier to pose the question whether every unitarizable group is amenable, which is still open today. On the other hand, the first examples of non-unitarizable groups were found by Ehrenpreis and Mautner [EM55], who showed that $\text{SL}_2(\mathbb{R})$ is not unitarizable. For discrete groups, explicit families of uniformly bounded non-unitarizable representations of free groups were later constructed by Mantero–Zappa [MZ83-1, MZ83-2], Pitlyk–Szwarc [PS86], and Bozejko [Bo87]. We refer the reader to [Pi01, Pi05, Oz06] for details. Since the unitarizability passes to subgroups, it follows that all discrete groups containing free subgroups are non-unitarizable. We also remark that examples of non-unitarizable groups not containing non-abelian free subgroups were given by Epstein–Monod [EM09], Osin [Os09] and Monod–Ozawa [MO10]. Therefore, there exist many uniformly bounded representations which are not unitarizable. The present article gives a new and unexpected connection between the study of unitarizability of groups and the understanding of the class $\mathcal{U}_{\text{fin}}$.

Finally, let us explain the idea of the proof. Theorem 1.2 (a) is shown in Theorem 2.7. For (b), (i) $\Leftrightarrow$ (ii) is Theorem 2.8 and (iii) $\Rightarrow$ (i) is Corollary 2.9. Finally, (i) $\Rightarrow$ (iii) is shown in Theorem 3.1. For the proof of (a) and (b) (i) $\Leftrightarrow$ (ii) and (iii) $\Rightarrow$ (i), we use our characterization [AM12-2] of the class $\mathcal{U}_{\text{fin}}$ via positive definite functions. The most difficult step is to prove (b) (i) $\Rightarrow$ (iii). Note that our situation is different from that of Vasilescu–Zsido [VZ74], where they showed, using the Ryll–Nardzewski Theorem, that if a uniformly bounded representation π generates a finite von Neumann algebra, then π is unitarizable. There does not seem to be an obvious way to deduce from the assumption that $G = H \rtimes_\pi \Gamma$ is of finite type the condition that $W^*(\pi(\Gamma))$ is a finite von Neumann algebra (observe that when π is an

irreducible unitary representation, then $H \rtimes_{\pi} \Gamma$ is of finite type, but $\pi(\Gamma)'' = \mathbb{B}(H)$ is not a finite von Neumann algebra if $\dim(H) = \infty$). Assume that $G = H \rtimes_{\pi} \Gamma$ is of finite type. Then there exists $f \in \mathcal{P}(H)$ which generates the neighborhood basis of 0 and satisfies $f(\pi(s)\xi) = f(\xi)$ ($s \in \Gamma, \xi \in H$). If $\dim(H) < \infty$, then by Bochner's Theorem, there exists a probability measure μ on H (regarded as a real Hilbert space) such that

$$f(\xi) = \int_H e^{ix \cdot \xi} d\mu(x), \quad \xi \in H.$$

If $\dim(H) = \infty$, Bochner's Theorem does not hold (e.g. the positive definite function $\xi \mapsto e^{-\|\xi\|^2}$ is *not* the Fourier transform of a probability measure on H). However, there exists a cylinder set probability measure μ defined on the algebraic dual H'_a of H , such that

$$(2) \quad f(\xi) = \int_{H'_a} e^{i\langle \chi, \xi \rangle} d\mu(\chi), \quad \xi \in H.$$

One can in fact show that μ is supported on a Hilbert-Schmidt extension of H . See [Ya85] for details. Intuitively, one may hope that π is unitarizable by the following argument: define

$$\langle \xi, \eta \rangle'_{\pi} := \int_{H'_a} \langle \chi, \xi \rangle \langle \chi, \eta \rangle d\mu(\chi), \quad \xi, \eta \in H,$$

which is (if well-defined) an $\mathbb{R}$ -bilinear form. Since f is π -invariant, its dual action $\tilde{\pi}$ of Γ on H'_a preserves μ , which then implies that $\langle \cdot, \cdot \rangle'_{\pi}$ is Γ -invariant: $\langle \pi(s)\xi, \pi(s)\eta \rangle'_{\pi} = \langle \xi, \eta \rangle'_{\pi}$ ($\xi, \eta \in H, s \in \Gamma$). By complexification, we get a $\pi(\Gamma)$ -invariant inner product $\langle \cdot, \cdot \rangle_{\pi}$. We may then hope that the inner product gives the same topology as the original inner product, which would imply that π is unitarizable (see Lemma 3.2). There are two difficulties in this approach: (a) we must guarantee that second moments exist:

$$\int_{H'_a} \langle \chi, \xi \rangle^2 d\mu(\chi) \stackrel{?}{<} \infty$$

and (b) we must prove that the new inner product is equivalent to the original one. It does not seem likely that these conditions hold in general. We remedy this by noncommutative integration theory: assume that $\alpha: G \rightarrow \mathcal{U}(M)$ is an embedding into the unitary group of a II_1 -factor M . Then by setting $\alpha(s) := \alpha(0, s)$ and $\alpha(\xi) := \alpha(\xi, 1)$ ($\xi \in H, s \in \Gamma$), we get $\alpha(s)\alpha(\xi)\alpha(s)^* = \alpha(\pi(s)\xi)$. Let $\widetilde{M}$ be the space of all unbounded closed operators affiliated with M . Since for each $\xi \in H$, $t \mapsto \alpha(t\xi)$ is a strongly continuous unitary group, there exists a self-adjoint operator $T(\xi) \in \widetilde{M}_{\text{sa}}$ such that $\alpha(t\xi) = e^{itT(\xi)}$ ($t \in \mathbb{R}$). One can show that $T: H \rightarrow \widetilde{M}_{\text{sa}}$ is an $\mathbb{R}$ -linear homeomorphism onto its range. If we can show that $T(\xi) \in L^2(M, \tau)$ for every $\xi \in H$ (let us call it the L^2 -condition), then

$$(3) \quad \langle \xi, \eta \rangle'_{\pi} := \langle T(\xi), T(\eta) \rangle_{L^2(M, \tau)}, \quad (\xi, \eta \in H)$$

is a well-defined inner-product, which is Γ -invariant by $\alpha(s)T(\xi)\alpha(s)^* = T(\pi(s)\xi)$ ($\xi \in H, s \in \Gamma$). Moreover, one can show that the induced norm is equivalent to the original norm. By complexification, we are done by Lemma 3.2. Again, the real question is whether the L^2 -condition holds in general, a question essentially equivalent to the existence of second moments of μ . However, since the von Neumann algebra $\mathcal{A} = \{\alpha(\xi); \xi \in H\}''$ generated by the image of α is commutative, we may represent it as $\mathcal{A} = L^{\infty}(X, m)$ for some probability space (X, m) equipped with an m -preserving Γ -action β such that the W^* -dynamical system $(\tilde{\mathcal{A}}, \text{Ad}(\alpha(\cdot)))$ is equivariantly $*$ -isomorphic to $(L^0(X, m), \beta)$. Here, $L^0(X, m)$ is the space of all measurable

maps on a probability space (X, m) equipped with the topology of convergence in measure. Since H is a Hilbert space (hence of Rademacher type 2, see [AK06, §6]), it is known (this is the so-called Maurey–Nikishin factorization [Ni70, Ma72]) that any continuous linear operator $T: H \rightarrow L^0(X, m) = \tilde{A}$ factors through a Hilbert space. That is, there exists a bounded operator $\tilde{T}: H \rightarrow L^2(X, m)$ and an m -a.e. positive function $\psi \in L^\infty(X, m)$ such that the following diagram commutes:

$$\begin{array}{ccc} H & \xrightarrow{T} & L^0(X, m) \\ & \searrow \tilde{T} & \swarrow M_\psi \\ & L^2(X, m) & \end{array} \quad \circ$$

Here, M_ψ is the multiplication operator by the function ψ . More precisely, there exists an m -a.e. positive $\varphi \in L^\infty(X, m)$ such that $\int_X \varphi(x) [T\xi](x)^2 dm(x) < \infty$ for every $\xi \in H$. Then $\tilde{T}\xi := M_{\varphi^{\frac{1}{2}}} \cdot T\xi$ and $\psi = \varphi^{-\frac{1}{2}}$ gives the factorization. This result was essentially proved first by Nikishin [Ni70]. Nikishin's result was later simplified by Maurey [Ma72] by using the theory of Rademacher types. We refer the reader to [Pi86, AK06, GR85] for more details. In §3, we show an equivariant version of the Maurey–Nikishin factorization, namely we use a convexity argument to ensure (Theorem 3.5) that the operator $\tilde{T}$ can be made Γ -equivariant if T is assumed to be Γ -equivariant. Then we can define $\langle \cdot, \cdot \rangle'_\pi$ by the formula (3) with T replaced by $\tilde{T}$. Since T is a real linear homeomorphism onto its range we can also show that $\tilde{T}$ is a real linear homeomorphism onto its range too. The last step is to show that on the image $\tilde{T}(H)$ of H under $\tilde{T}$, the topology of the convergence in measure agrees with the L^2 -topology (Lemma 3.13). This is somewhat surprising: note that these two topologies do not agree on the whole of $L^2(X, m)$. The proof is done by using the fact [Be10, AM12-1] that on $\tilde{M}$, the strong resolvent topology and the τ -measure topology of Nelson [Ne74] agree. Once this is done, it is straightforward to show that $\langle \cdot, \cdot \rangle'_\pi$ generates the same topology as the original inner product, and therefore π is unitarizable.

Finally, let us point out that as a byproduct of the existence of an equivariant Maurey–Nikishin factorization, we find a new and useful description of positive definite functions on a Hilbert space (compare (2) with (4)) (see also the comments after Corollary 3.16):

Theorem 1.3 (Corollary 3.16). *Let H be a separable real Hilbert space, and let $f \in \mathcal{P}(H)$. Then there exist*

- (i) *a compact metrizable space X and a Borel probability measure m on X ,*
- (ii) *a bounded linear operator $T: H \rightarrow L^2(X, m; \mathbb{R})$, and*
- (iii) *a positive measurable function $\varphi \in L^\infty(X, m)$,*

such that

$$(4) \quad f(\xi) = \int_X e^{i\varphi(x)[T\xi](x)} dm(x), \quad \xi \in H.$$

Moreover:

- (iv) *If f generates a neighborhood basis of 0, then T can be chosen to be a real-linear homeomorphism onto its range.*
- (v) *We may arrange T and φ so that if $\mathcal{G} = \{u \in O(H); f(u\xi) = f(\xi) \ (\xi \in H)\}$ is the subgroup of the orthogonal group of H fixing f pointwise, then there exists a continuous*

homomorphism β from $\mathcal{G}$ to the group $\text{Aut}(X, m)$ of all m -preserving automorphisms of X with the weak topology, such that

$$[T(u\xi)](x) = \beta(u)[T(\xi)](x), \quad \varphi(\beta(u)x) = \varphi(x), \quad u \in \mathcal{G}, \xi \in H, \text{ } m\text{-a.e. } x \in X.$$

2. THE CLASS $\mathcal{U}_{\text{fin}}$ AND UNIFORMLY BOUNDED REPRESENTATIONS

In this section we recall Popa's class $\mathcal{U}_{\text{fin}}$ of finite type Polish groups and consider when a specific semidirect group belongs to this class. The results of this section are extracted from the first named author's PhD thesis, jointly proved with the second named author.

Definition 2.1. Let G, G' be topological groups.

- (i) We say that G is *embeddable* into G' , if there exists a topological group isomorphism π from G onto a closed subgroup of G' . Such π is called an *embedding*.
- (ii) We denote by $\mathcal{P}(G)$ the set of all normalized continuous positive definite functions on G , and by $\mathcal{P}(G)_{\text{inv}}$ the set of all $f \in \mathcal{P}(G)$ which is conjugation-invariant, i.e., $f(g^{-1}xg) = f(x)$ holds for all $x, g \in G$. An element of $\mathcal{P}(G)_{\text{inv}}$ is called a *character* on G .

Definition 2.2 ([Po07]). A Polish group G is of *finite type* if G is embeddable into the unitary group $\mathcal{U}(M)$ of some finite von Neumann algebra M equipped with strong operator topology. The class of finite type Polish groups is denoted by $\mathcal{U}_{\text{fin}}$.

If G is a finite type Polish group, then by [Po07, Lemma 2.6], the above M may chosen to be a type II_1 -factor with separable predual.

Definition 2.3. Let G be a topological group.

- (i) G is called *unitarily representable* if G is embeddable into $\mathcal{U}(\ell^2(\mathbb{N}))$ equipped with strong operator topology.
- (ii) G is called *SIN* (small invariant neighborhood) if for any neighborhood V of G at the identity e , there exists a neighborhood $U \subset V$ of e which is conjugation invariant, i.e., $g^{-1}Ug = U$, $\forall g \in G$. Such U is called an *invariant neighborhood* of G .

If G is metrizable, then G is SIN if and only if G admits a bi-invariant metric d which is compatible with its topology.

For more details about unitary representability, see e.g., [Gal09, Gao05, Me02]. We will use the following characterization of the finite type Polish groups (cf. [Gao05]).

Theorem 2.4 ([AM12-2]). *For a Polish group G , the following conditions are equivalent.*

- (i) G is of finite type.
- (ii) There exists a family $\mathcal{F} \subset \mathcal{P}(G)_{\text{inv}}$ which generates a neighborhood basis of the identity e_G of G . That is, for each neighborhood V at the identity, there are functions $f_1, \dots, f_n \in \mathcal{F}$ and open sets $\mathcal{O}_1, \dots, \mathcal{O}_n$ in $\mathbb{C}$ such that

$$e_G \in \bigcap_{i=1}^n f_i^{-1}(\mathcal{O}_i) \subset V.$$

- (iii) There exists $f \in \mathcal{P}(G)_{\text{inv}}$ which generates a neighborhood basis of the identity of G .

- (iv) *There exists a family $\mathcal{F} \subset \mathcal{P}(G)_{\text{inv}}$ which separates the identity of G and closed subsets A with $A \not\ni e_G$. That is, for each closed subset A with $A \not\ni e_G$, there exists $f \in \mathcal{F}$ such that*

$$\sup_{x \in A} |f(x)| < |f(e_G)|.$$

- (v) *There exists $f \in \mathcal{P}(G)_{\text{inv}}$ which separates the identity of G and closed subsets A with $A \not\ni e_G$.*

We start from the following characterization for the semidirect product of a SIN group by a SIN group to be SIN. Denote by $\mathcal{N}(G)$ the set of all neighborhoods of identity of a topological group G .

Theorem 2.5. *Let N, K be SIN-groups. Let $\alpha: K \rightarrow \text{Aut}(N)$ be a continuous action of K on N . Then the following two conditions are necessary and sufficient for $N \rtimes_{\alpha} K$ to be a SIN-group.*

- (SIN. 1) *For any $U \in \mathcal{N}(N)$, there exist $V \in \mathcal{N}(N)$ with $V \subset U$ such that*

$$\alpha_k(V) = V, \quad k \in K.$$

- (SIN. 2) *The action α is bounded. That is, for any $V \in \mathcal{N}(N)$, there exists $W \in \mathcal{N}(K)$ such that*

$$n^{-1}\alpha_k(n) \in V, \quad n \in N, \quad k \in W.$$

Proof. We denote by e_N (resp. e_K) the unit of N (resp. K).

We first prove sufficiency of the two conditions. Fix an arbitrary $U \in \mathcal{N}(N \rtimes_{\alpha} K)$. By definition of the product topology, there exist $V'' \in \mathcal{N}(N)$ and $W' \in \mathcal{N}(K)$ such that $(e_N, e_K) \in V'' \times W' \subset U$ holds. Since N is SIN, there exists an invariant neighborhood $V' \in \mathcal{N}(N)$ which satisfies $V' \cdot V' \subset V''$. By (SIN. 1), we may take $V \in \mathcal{N}(N)$ such that $V \subset V'$ and

$$\alpha_k(V) = V, \quad k \in K.$$

By (SIN. 2) and the SIN property of K , there exists an invariant neighborhood $W \in \mathcal{N}(K)$ satisfying $W \subset W'$ and

$$n^{-1}\alpha_k(n) \in V, \quad n \in N, \quad k \in W.$$

Indeed, by (SIN. 2) there is $W_0 \in \mathcal{N}(K)$ such that $n^{-1}\alpha_k(n) \in V$ holds for all $n \in N$ and $k \in W_0$. By the SIN property of K , there is an invariant neighborhood $W \in \mathcal{N}(K)$ contained in $W_0 \cap W'$. It is then clear that this W satisfies the above requirement.

Let $(n_1, k_1) \in V \times W$ and $(n_2, k_2) \in N \rtimes_{\alpha} K$. We see that

$$\begin{aligned} (n_2, k_2)(n_1, k_1)(n_2, k_2)^{-1} &= (n_2\alpha_{k_2}(n_1), k_2k_1)(\alpha_{k_2^{-1}}(n_2^{-1}), k_2^{-1}) \\ &= (n_2\alpha_{k_2}(n_1)\alpha_{k_2k_1k_2^{-1}}(n_2^{-1}), k_2k_1k_2^{-1}) \\ &= \left(\{n_2\alpha_{k_2}(n_1)n_2^{-1}\} \{n_2\alpha_{k_2k_1k_2^{-1}}(n_2^{-1})\}, k_2k_1k_2^{-1} \right) \\ &\in (n_2Vn_2^{-1} \cdot V) \times W \\ &\subset (n_2V'n_2^{-1} \cdot V') \times W' = (V' \cdot V') \times W' \\ &\subset U. \end{aligned}$$

Hence, we have $g^{-1}(V \times W)g \subset U$ for all $g \in N \rtimes_{\alpha} K$. This shows

$$U' := \bigcup_{g \in N \rtimes_{\alpha} K} g^{-1}(V \times W)g \subset U$$

is an invariant neighborhood of $e_{N \rtimes_\alpha H}$ contained in U . Since U is arbitrary, this shows $N \rtimes_\alpha K$ is a SIN-group.

We now prove necessity of the two conditions. Suppose $N \rtimes_\alpha K$ is a SIN-group. Given $U \in \mathcal{N}(N)$. Then $U \times K \in \mathcal{N}(N \rtimes_\alpha K)$. Thus by SIN-property of $N \rtimes_\alpha K$ we find an invariant neighborhood $A \in \mathcal{N}(N \rtimes_\alpha K)$ contained in $U \times K$. Then we may find $U_0 \in \mathcal{N}(N)$ and $W \in \mathcal{N}(K)$ such that $U_0 \times W \subset A$. Since A is invariant, we have

$$g(U_0 \times \{e_K\})g^{-1} \subset A, \quad g \in N \rtimes_\alpha K.$$

Inparticular, for all $k \in K$ and $n \in U_0$, we have

$$(e_N, k)(n, e_K)(e_N, k)^{-1} = (\alpha_k(n), e_K) \in A \subset U \times K.$$

This shows $\alpha_k(n) \in U$ for all $k \in K$ and $n \in U_0 \subset U$. Then

$$V := \bigcup_{k \in K} \alpha_k(U_0) \subset U$$

is a neighborhood of e_N satisfying $\alpha_k(V) = V$ for all $k \in K$. Hence we conclude that (SIN. 1) holds.

In order to prove (SIN. 2), suppose $V \in \mathcal{N}(N)$ is given. Again by the same argument as above, we may find an invariant neighborhood $A \in \mathcal{N}(N \rtimes_\alpha K)$ contained in $V \times K$ and $V_0 \in \mathcal{N}(N)$ and $W \in \mathcal{N}(K)$ with $V_0 \times W \subset A \subset V \times K$. Since A is invariant, we have in particular

$$g(\{e_N\} \times W)g^{-1} \subset A \subset V \times K, \quad g \in N \rtimes_\alpha K.$$

Hence for $n \in N$ and $k \in W$, we have

$$(n, e_K)(e_N, k)(n, e_K)^{-1} = (n\alpha_k(n^{-1}), k) \in A \subset V \times K.$$

This shows $n\alpha_k(n^{-1}) \in V$ for all $k \in W$. This proves (SIN. 2). $\square$

Remark 2.6. A semidirect product $G \rtimes_\alpha \Gamma$ of a SIN group G by a discrete group Γ always satisfies (SIN. 2) (take $W = \{e_K\}$).

As an immediate corollary to Theorem 2.5, we obtain:

Theorem 2.7. *Let Γ be a countable discrete group, H be a separable Hilbert space and $\pi: \Gamma \rightarrow \text{GL}(H)$ be an invertible representation of Γ . Then the semidirect product group $G := H \rtimes_\pi \Gamma$ is unitarily representable. Moreover, the following conditions are equivalent.*

- (1) G is SIN.
- (2) π is uniformly bounded, i.e., $\|\pi\| := \sup_{s \in \Gamma} \|\pi(s)\| < \infty$.

Proof. The unitarily representable group H has a positive definite function $f(\xi) := e^{-\|\xi\|^2}$, $\xi \in H$ and identifying H with $H \times \{e\} \subset H \rtimes_\pi \Gamma$, we may extend f to be a continuous positive definite function on $H \rtimes_\pi \Gamma$ by taking value 0 outside $H \times \{e\}$. It is immediate to see that this extended function still generates the neighborhood basis of the identity. Then by [Gao05, Theorem 2.1], G is unitarily representable.

(1) $\Rightarrow$ (2) Assume that G is a unitarily representable and SIN group. Let $B_\delta(0)$ be the open δ -neighborhood of H at 0 ($\delta > 0$). Then by (SIN. 1), there exists a neighborhood $V \subset B_\delta(0)$ of H at 0 such that $\pi(s)V = V$, ($s \in \Gamma$). Since V is a neighborhood, there exists $\varepsilon > 0$ such that $B_\varepsilon(0) \subset V$ holds. Thus $\pi(s)B_\varepsilon(0) \subset B_\delta(0)$ holds for all $s \in \Gamma$. This shows that $\|\pi(s)\| \leq \frac{\delta}{\varepsilon}$ for every $s \in \Gamma$. Therefore π is uniformly bounded.

(2) $\Rightarrow$ (1) Assume that π is uniformly bounded. To prove that G is SIN, by Theorem 2.5 we have

only to check (SIN.1) (see Remark 2.6). Consider $B_\varepsilon(0)$ ($\varepsilon > 0$), and $C := 1 + \sup_{s \in \Gamma} \|\pi(s)\| < \infty$. Then for all $\xi \in B_{\varepsilon/C}(0)$, $s \in \Gamma$, we have

$$\|\pi(s)\xi\| \leq C\|\xi\| < \varepsilon.$$

This implies $\pi(s)B_{\varepsilon/C}(0) \subset B_\varepsilon(0)$ for all $s \in \Gamma$. Then $V := \bigcup_{s \in \Gamma} \pi(s)B_{\varepsilon/C}(0)$ satisfies (SIN.1). $\square$

Theorem 2.8. *Let Γ be a countable discrete group, H be a separable Hilbert space and $\pi : \Gamma \rightarrow \mathbb{B}(H)$ be a uniformly bounded representation of Γ . Then $H \rtimes_\pi \Gamma$ is of finite type if and only if there exists $f \in \mathcal{P}(H)$ such that*

$$(F.1) \quad f(\pi(s)\xi) = f(\xi), \quad \forall s \in \Gamma, \xi \in H,$$

$$(F.2) \quad f \text{ generates a neighborhood basis of } 0 \in H.$$

Proof. We first prove “if” part. For this, we assume that there exists a positive, continuous positive definite function f on H which satisfies (F.1) and (F.2). Then one can extend f on $H \rtimes_\pi \Gamma$ by

$$\tilde{f}(\xi, s) := \begin{cases} f(\xi) & (s = 1), \\ 0 & (s \neq 1). \end{cases}$$

It is easy to see that $\tilde{f}$ is continuous, positive definite and generates a neighborhood basis of the identity. Take arbitrary $\xi, \eta \in H$, $s, t \in \Gamma$. Since

$$(\eta, t)(\xi, s)(\eta, t)^{-1} = (\eta + \pi(t)\xi - \pi(tst^{-1})\eta, tst^{-1}),$$

we have

$$\begin{aligned} \tilde{f}((\eta, t)(\xi, s)(\eta, t)^{-1}) &= \begin{cases} \tilde{f}(\pi(t)\xi, 1) & (s = 1), \\ 0 & (s \neq 1). \end{cases} \\ &= \begin{cases} f(\xi) & (s = 1), \\ 0 & (s \neq 1). \end{cases} \\ &= \tilde{f}(\xi, s). \end{aligned}$$

Hence $H \rtimes_\pi \Gamma$ is of finite type.

Conversely, let us assume that $H \rtimes_\pi \Gamma$ is of finite type. Then there exists a positive, continuous positive definite class function $\tilde{f}$ on $H \rtimes_\pi \Gamma$ which generates a neighborhood basis of the identity. We define a positive definite function f on H by

$$f(\xi) := \tilde{f}(\xi, 1), \quad \xi \in H.$$

It is easy to check that f is positive, continuous and satisfies (F.2). Since

$$(\eta, t)(\xi, 1)(\eta, t)^{-1} = (\pi(t)\xi, 1)$$

and $\tilde{f}$ is a class function, we obtain

$$\begin{aligned} f(\xi) &= \tilde{f}(\xi, 1) = \tilde{f}((\eta, t)(\xi, 1)(\eta, t)^{-1}) \\ &= \tilde{f}(\pi(t)\xi, 1) = f(\pi(t)\xi) \end{aligned}$$

so that f satisfies (F.1). $\square$

Corollary 2.9. *Let Γ be a countable discrete group, H be a separable Hilbert space and $\pi : \Gamma \rightarrow \mathbb{B}(H)$ be a uniformly bounded representation of Γ . If π is unitarizable, then $H \rtimes_\pi \Gamma$ is of finite type.*

Proof. Since π is unitarizable, there exists an invertible operator $T \in \mathbb{B}(H)$ such that $T\pi(s)T^{-1}$ is unitary for every $s \in \Gamma$. Define a positive definite function f on H by

$$f(\xi) := e^{-\|T\xi\|^2}, \quad \xi \in H.$$

Then f is positive, continuous and satisfies (F.2). Moreover

$$f(\pi(s)\xi) = e^{-\|T\pi(s)\xi\|^2} = e^{-\|\{T\pi(s)T^{-1}\}T\xi\|^2} = e^{-\|T\xi\|^2} = f(\xi)$$

so that (F.1) holds. $\square$

Since amenable groups are unitarizable ([Da50, Di50, NT51]), we obtain:

Corollary 2.10. *Let Γ be a countable discrete group, H be a separable Hilbert space and $\pi : \Gamma \rightarrow \mathbb{B}(H)$ be a uniformly bounded representation of Γ . Assume that Γ is amenable. Then $H \rtimes_{\pi} \Gamma$ is of finite type.*

Remark 2.11. We remark that if Γ is amenable, then so is $H \rtimes_{\pi} \Gamma$, so that the conclusion of Corollary 2.10 also follows from the fact [AM12-2, Theorem 2.2] that an amenable unitarily representable SIN group is of finite type.

The purpose of the next section is to prove a converse to Corollary 2.9.

3. EQUIVARIANT MAUREY–NIKISHIN FACTORIZATION

In this section, we prove the last and crucial step, Theorem 1.2 (b) (i) $\Rightarrow$ (iii). Let $\pi : \Gamma \rightarrow H$ be a uniformly bounded representation of a discrete countable group Γ on a separable Hilbert space H , as before.

Theorem 3.1. *If $G = H \rtimes_{\pi} \Gamma$ is of finite type, then π is unitarizable.*

Throughout this section, we assume that $G = H \rtimes_{\pi} \Gamma$ is of finite type. Our aim is to find a Γ -invariant inner product and apply the following classical result due to Dixmier:

Lemma 3.2 (Dixmier). *Let $\pi : \Gamma \rightarrow \mathbb{B}(H)$ be a uniformly bounded representation. Then π is unitarizable if and only if there exists an inner product $\langle \cdot, \cdot \rangle'$ on H which generates the same Hilbert space topology and which is π -invariant, i.e., $\langle \pi(s)\xi, \pi(s)\eta \rangle' = \langle \xi, \eta \rangle'$ ($\xi, \eta \in H, s \in \Gamma$) holds.*

By assumption, there exists an embedding (as a topological group) $\alpha : G \rightarrow \mathcal{U}(M)$, where M is some II_1 -factor with separable predual and with the normal faithful tracial state τ . We let M act on $L^2(M, \tau)$. As usual, we identify $H \subset G \subset \mathcal{U}(M)$ and $\Gamma \subset \mathcal{U}(M)$. That is, $\xi \in H$ is identified with $\alpha(\xi) = \alpha(\xi, 0)$, and $s \in \Gamma$ is identified with $\alpha(s) = \alpha(0, s)$. Therefore under this identification, the uniformly bounded G -action π on H is implemented by unitaries in M :

$$(5) \quad \alpha(\pi(s)\xi) = \alpha((0, s)(\xi, 1)(0, s^{-1})) = \alpha(s)\alpha(\xi)\alpha(s)^*.$$

Now, let $\widetilde{M}$ be the space of all (possibly unbounded) closed and densely defined operators T on $L^2(M, \tau)$ which are affiliated with M . There is a completely metrizable topology on $\widetilde{M}$, called the τ -measure topology [Ne74]. A neighborhood basis of 0 in the τ -measure topology is given by the family $\{N(\varepsilon, \delta); \varepsilon, \delta > 0\}$, where

$$N(\varepsilon, \delta) := \{A \in \widetilde{M}; \exists p \in \text{Proj}(M), \tau(p^{\perp}) < \delta, \|Ap^{\perp}\| < \varepsilon\},$$

where $\text{Proj}(M)$ is the lattice of all projections of M . In [AM12-1], we showed that the τ -measure topology coincides with the strong resolvent topology (SRT) (the same result is obtained by Beltita [Be10] using the standard form). In particular, if $A_n, A \in \widetilde{M}$ are self-adjoint, then $A_n \xrightarrow{n \rightarrow \infty} A$ in the τ -measure topology, if and only if $\|(A_n - i)^{-1} - (A - i)^{-1}\|_2 \xrightarrow{n \rightarrow \infty} 0$, if and only if (by the Kato-Trotter Theorem) $\|e^{itA_n} - e^{itA}\|_2 \xrightarrow{n \rightarrow \infty} 0$ for every $t \in \mathbb{R}$ (in this case the convergence is compact uniform in t). As was shown by Murray-von Neumann, the space $\widetilde{M}$, although not locally convex, is actually a topological $*$ -algebra with respect to the τ -measure topology. The addition (resp. the multiplication) is defined by $(A, B) \mapsto A + \overline{B}$ (resp. $(A, B) \mapsto \overline{AB}$), where the bar denotes the operator closure (it asserts that $A + B$, AB are always closable and in fact $\overline{A + B}$ is self-adjoint whenever A, B are).

Now let $\xi \in H$. Then since α is continuous, $\mathbb{R} \ni t \mapsto \alpha(t\xi) \in \mathcal{U}(M)$ is a strongly continuous one-parameter unitary group. Thus by Stone's Theorem, there exists a unique self-adjoint operator $T(\xi) \in \widetilde{M}$ such that

$$\alpha(t\xi) = e^{itT(\xi)}, \quad t \in \mathbb{R}, \xi \in H.$$

Proposition 3.3. *The map $T: H \rightarrow \widetilde{M}_{\text{sa}}$ is a real-linear homeomorphism onto its range (with respect to the τ -measure topology or the SRT). Moreover, the following covariance property holds.*

$$(6) \quad T(\pi(s)\xi) = \alpha(s)T(\xi)\alpha(s)^*, \quad s \in \Gamma, \xi \in H.$$

Proof. Let $s \in \Gamma$ and $\xi_n, \xi, \eta \in H$ ($n \in \mathbb{N}$). Since α is a topological isomorphism onto its range,

$$\begin{aligned} \xi_n - \xi \xrightarrow{n \rightarrow \infty} 0 &\Leftrightarrow \alpha(t\xi_n) - \alpha(t\xi) = e^{itT(\xi_n)} - e^{itT(\xi)} \xrightarrow{n \rightarrow \infty} 0 \text{ for all } t \in \mathbb{R} \text{ (SOT)} \\ &\Leftrightarrow T(\xi_n) \xrightarrow{n \rightarrow \infty} T(\xi) \text{ (SRT)}. \end{aligned}$$

Thus T is a homeomorphism from H onto its range. Also, for all $s, t \in \mathbb{R}$, we have

$$\alpha(ts\xi) = e^{itsT(\xi)} = \alpha(t(s\xi)) = e^{itT(s\xi)}.$$

Since t is arbitrary, the uniqueness part of Stone's Theorem asserts that $T(s\xi) = sT(\xi)$. Now since H is a commutative (additive) group, all $\{T(\xi); \xi \in H\}$ are strongly commuting self-adjoint operators. Thus by Kato-Trotter Theorem,

$$\begin{aligned} e^{it\overline{T(\xi) + T(\eta)}} &= \text{SOT} - \lim_{n \rightarrow \infty} (e^{it\frac{1}{n}T(\xi)} e^{it\frac{1}{n}T(\eta)})^{\frac{1}{n}} = e^{itT(\xi)} e^{itT(\eta)} \\ &= \alpha(t\xi)\alpha(t\eta) = \alpha(t(\xi + \eta)) \\ &= e^{itT(\xi + \eta)}. \end{aligned}$$

Therefore we obtain $T(\xi + \eta) = \overline{T(\xi) + T(\eta)}$. Therefore T is a real-linear homeomorphism onto its range. Finally, by the unitary covariance of the Borel functional calculus,

$$\begin{aligned} e^{itT(\pi(s)\xi)} &= \alpha(t\pi(s)\xi) = \alpha(s)\alpha(t\xi)\alpha(s)^* \\ &= \alpha(s)e^{itT(\xi)}\alpha(s)^* \\ &= e^{it\alpha(s)T(\xi)\alpha(s)^*}. \end{aligned}$$

Therefore $T(\pi(s)\xi) = \alpha(s)T(\xi)\alpha(s)^*$. □

Let $\mathcal{A} := \{\alpha(\xi); \xi \in H\}''$ be the commutative von Neumann subalgebra of M generated by the image of α . By Proposition 3.18, there exists a probability space (X, m) such that $(\mathcal{A}, \tau|_{\mathcal{A}}) \cong (L^\infty(X, m), \int_X \cdot dm)$. Then by (6), the formula

$$(7) \quad \beta_s := \text{Ad}(\alpha(s))|_{\mathcal{A}}, \quad s \in \Gamma.$$

defines a trace-preserving action of Γ on $\mathcal{A} = L^\infty(X, m)$. This action also defines a probability measure preserving action on X . We denote this action by $\Gamma \times X \ni (s, x) \mapsto s \cdot x \in X$. Therefore, $\beta_s(\varphi) = \varphi \circ s^{-1}$, $\varphi \in L^\infty(X, m)$. Below we keep this setting.

Definition 3.4. We denote by $L^0(X, m)$ the space of all m -measurable $\mathbb{C}$ -valued functions on X , and $L^0(X, m)_+$ its positive cone. We also define the Γ -action on $L^0(X, m)$ by $\beta_s = \text{Ad}(\alpha(s))|_{L^0(X, m)}$ ($s \in \Gamma$) (we use the same symbol β_s as the Γ -action on $L^\infty(X, m)$, but there would be no danger of confusion).

Equipped with the convergence in measure, $L^0(X, m)$ is a (non-locally convex) completely metrizable topological vector space. Note that under the identification $\mathcal{A} = L^\infty(X, m)$, we obtain $\tilde{\mathcal{A}} = L^0(X, m) \subset \tilde{M}_{\text{sa}}$ (cf. [AM12-1]). What we will do is to construct a Γ -equivariant Maurey–Nikishin factorization of the map $T: H \rightarrow L^0(X, m) \subset \tilde{M}_{\text{sa}}$. We almost exactly follow Maurey’s argument as in [GR85, Chapter VI.3], but we also need to take care of the Γ -action, which makes it necessary to refine some of the arguments.

Theorem 3.5 (Equivariant Maurey–Nikishin Factorization). *The map $T: H \rightarrow L^0(X, m)$ factors Γ -equivariantly through a Hilbert space. More precisely, there exists a Γ -invariant function $\varphi \in L^0(X, m)$ which is m -a.e. positive and*

$$(8) \quad \int_X \varphi(x) [T\xi](x)^2 dm(x) \leq \|\xi\|^2, \quad \xi \in H.$$

In particular, the following diagram commutes:

$$\begin{array}{ccc} H & \xrightarrow{T} & L^0(X, m) \\ & \searrow \tilde{T} & \nearrow M_\psi \\ & L^2(X, m) & \end{array} \quad \circlearrowright$$

Here, M_ψ is the multiplication operator by the Γ -invariant function $\psi = \varphi^{-\frac{1}{2}}$ and $\tilde{T}\xi := M_{\varphi^{\frac{1}{2}}} \cdot T\xi$ is Γ -equivariant.

Definition 3.6. For $\varphi \in L^0(X, m)$, we define its *non-increasing rearrangement* $\varphi^*: [0, 1] \rightarrow [0, \infty)$ by

$$\varphi^*(t) := \inf\{\lambda > 0; m(\{x \in X; |\varphi(x)| > \lambda\}) \leq t\}, \quad t \in [0, 1].$$

Then clearly $(c\varphi)^*(t) = c\varphi^*(t)$ ($c > 0$, $t \geq 0$) holds.

The next lemmata are elementary.

Lemma 3.7. *Let $S: H \rightarrow L^0(X, m)$ be a linear operator. Then the following conditions are equivalent.*

- (i) S is continuous.
- (ii) There exists a non-increasing map $C: (0, \infty) \rightarrow [0, 1]$ such that $\lim_{\lambda \rightarrow \infty} C(\lambda) = 0$ and

$$(9) \quad m(\{x \in X; |S\xi(x)| \geq \lambda\|\xi\|\}) \leq C(\lambda) \quad (\lambda > 0, \xi \in H).$$

(iii) For every $t > 0$, there exists $K(t) > 0$ such that

$$(10) \quad (S\xi)^*(t) \leq K(t)\|\xi\| \quad (\xi \in H).$$

Proof. (i) $\Rightarrow$ (ii) For each $\varepsilon, \delta > 0$, the set $U(\varepsilon, \delta) := \{f \in L^0(X, m); m(\{x \in X; |f(x)| \geq \varepsilon\}) \leq \delta\}$ is an neighborhood of 0 in the measure topology. Fix $\lambda > 0$. Then by the continuity of S , for each $\delta > 0$, $S^{-1}(U(\lambda, \delta))$ is an neighborhood of 0 in H , so that there exists $r_{\delta, \lambda} > 0$ such that $S\xi \in U(\lambda, \delta)$ holds for every $\xi \in H$ with $\|\xi\| \leq r_{\delta, \lambda}$. This implies that the implication $\|\xi\| \leq 1 \Rightarrow S\xi \in U(\lambda, \delta/r_{\delta, \lambda})$ holds. Define

$$C(\lambda) := \inf\{\delta > 0; S\xi \in U(\lambda, \delta) \text{ for all } \xi \in H \text{ with } \|\xi\| \leq 1\}.$$

It is then clear that $C(\cdot)$ is nonincreasing and (9) holds. We claim that $\lim_{\lambda \rightarrow \infty} C(\lambda) = 0$. Assume by contradiction that there exists $\delta > 0$ such that $C(\lambda) \geq \delta$ holds for all $\lambda > 0$. Then for each $n \in \mathbb{N}$, there exists $\xi_n \in H$ with $\|\xi_n\| \leq 1$, such that $S\xi_n \notin U(n, \delta - \frac{1}{n})$, i.e., $m(\{x \in X; |S\xi_n(x)| \geq n\}) > \delta - \frac{1}{n}$ holds. Then $\eta_n := \frac{1}{n}\xi_n$ ($n \in \mathbb{N}$) satisfies $\|\eta_n\| \xrightarrow{n \rightarrow \infty} 0$, but $m(\{x \in X; |S\eta_n(x)| \geq 1\}) > \delta - \frac{1}{n}$, so $S\eta_n \not\rightarrow 0$ in measure, a contradiction.

(ii) $\Rightarrow$ (iii) Let $t > 0$. Since $\lim_{\lambda \rightarrow \infty} C(\lambda) = 0$, there exists $K(t) > 0$ such that $C(\lambda) < t$ ($\lambda \geq K(t)$) holds. Then for $\xi \in H$ with $\|\xi\| \leq 1$, $m(\{x \in X; |S\xi(x)| > \lambda\}) \leq C(\lambda)$, so that $(S\xi)^*(t) \leq K(t)$. For general $\xi \neq 0$, we have $(S\|\xi\|^{-1}\xi)^*(t) \leq \|\xi\|^{-1}K(t)$, whence (10) holds.

(iii) $\Rightarrow$ (i) Since H and $L^0(X, m)$ are metrizable, it suffices to show the sequential continuity of S . Let $\xi_n \in H$ be such that $\|\xi_n\| \xrightarrow{n \rightarrow \infty} 0$. Let $\varepsilon, \delta > 0$. By assumption, we have $(S\xi_n)^*(\delta) \leq K(\delta)\|\xi_n\| \xrightarrow{n \rightarrow \infty} 0$. Therefore there exists $N \in \mathbb{N}$ such that $(S\xi_n)^*(\delta) < \varepsilon$ holds for all $n \geq N$. In particular, we have $m(\{x \in X; |S\xi_n(x)| \geq \varepsilon\}) \leq \delta$. Since $\varepsilon, \delta > 0$ are arbitrary, $S\xi_n$ converges to 0 in measure. $\square$

Lemma 3.8. *Let $\varphi \in L^0(X, m)$. Then for every $\delta > 0$, $m(\{x \in X; |\varphi(x)| > \varphi^*(\delta)\}) \leq \delta$ holds.*

Proof. Let $\lambda := \varphi^*(\delta)$. Then for each $n \in \mathbb{N}$, there exists $\lambda_n \in (\lambda, \lambda + \frac{1}{n})$ such that $\mu(\{x \in X; |\varphi(x)| > \lambda_n\}) \leq \delta$. Thus

$$\begin{aligned} m(\{x \in X; |\varphi(x)| > \lambda\}) &\leq \lim_{n \rightarrow \infty} m(\{x \in X; |\varphi(x)| > \lambda + \frac{1}{n}\}) \\ &\leq \limsup_{n \rightarrow \infty} m(\{x \in X; |\varphi(x)| > \lambda_n\}) \\ &\leq \delta. \end{aligned}$$

$\square$

The crucial step in our proof is to establish the following proposition. This is the step which allows to make the construction of the factorization canonical and hence equivariant with respect to a group of symmetries.

Proposition 3.9. *Let A be a convex subset of $L^0(X, m)_+$ which is globally Γ -invariant. Then for every $\varepsilon > 0$, there is a Γ -invariant measurable subset $S(\varepsilon) \subset X$ with $m(X \setminus S(\varepsilon)) \leq \varepsilon$ and*

$$(11) \quad \sup_{\varphi \in A} \int_{S(\varepsilon)} \varphi(x) dm(x) \leq 2 \sup_{\varphi \in A} \varphi^*\left(\frac{\varepsilon}{2}\right).$$

This is a Γ -invariant version of [GR85, Proposition 3.1]. In order to show Proposition 3.9, we use the following variant of the Minimax Lemma:

Lemma 3.10 (Minimax Lemma). *Let X, Y be topological vector spaces, and let A (resp. B) be a convex subset of X (resp. Y). Assume that B is compact. If the function $\Phi: A \times B \rightarrow \mathbb{R} \cup \{+\infty\}$ satisfies:*

- (i) $\Phi(\cdot, b)$ is a concave function on A for each $b \in B$,
- (ii) $\Phi(a, \cdot)$ is a convex function on B for each $a \in A$,
- (iii) $\Phi(a, \cdot)$ is lower-semicontinuous on B for each $a \in A$,

then the following identity holds:

$$\min_{b \in B} \sup_{a \in A} \Phi(a, b) = \sup_{a \in A} \min_{b \in B} \Phi(a, b).$$

Proof. See [GR85, Appendix A.2]. □

Proof of Proposition 3.9. Define

$$B := \left\{ \varphi \in L^0(X, m); 0 \leq \varphi \leq 1 \text{ (m-a.e.)}, \int_X \varphi(x) dm(x) \geq 1 - \frac{\varepsilon}{2} \right\}.$$

Then we first check that B is a globally Γ -invariant, weakly compact, convex subset of the Hilbert space $L^2(X, m)$: indeed, $B \subset L^\infty(X, m) \subset L^2(X, m)$ is clearly a convex set contained in the unit ball of $L^2(X, m)$ (any $\varphi \in B$ satisfies $\int_X |\varphi|^2 dm \leq \int_X 1 dm = 1$). Since $m \circ \beta_s = m$ ($s \in \Gamma$), the Γ -invariance is clear. Suppose $(\varphi_n)_{n=1}^\infty$ is a sequence in B converging weakly to $\varphi \in L^2(X, m)$. For each $k \in \mathbb{N}$, let $E_k = \{x \in X; \varphi(x) \leq -\frac{1}{k}\}$. Then

$$0 \leq \lim_{n \rightarrow \infty} \langle \varphi_n, 1_{E_k} \rangle_{L^2(X)} = \langle \varphi, 1_{E_k} \rangle_{L^2(X)} = \int_{E_k} \varphi dm \leq -\frac{1}{k} m(E_k) \leq 0,$$

so that $m(E_k) = 0$. Therefore $E = \{x \in X; \varphi(x) < 0\} = \bigcup_{k \in \mathbb{N}} E_k$ satisfies $m(E) = 0$, so $\varphi \geq 0$ a.e. Similarly, $\varphi \leq 1$ a.e. holds, and

$$\int_X \varphi dm = \langle \varphi, 1 \rangle_{L^2(X)} = \lim_{n \rightarrow \infty} \langle \varphi_n, 1 \rangle \geq 1 - \frac{\varepsilon}{2}.$$

Therefore $\varphi \in B$. This shows that B is weakly closed, so that B is a weakly compact convex subset of $L^2(X, m)$.

Next, define $\Phi: A \times B \rightarrow \mathbb{R} \cup \{+\infty\}$ by

$$\Phi(\varphi, \psi) := \int_X \psi \varphi dm, \quad (\psi \in A, \varphi \in B).$$

It is clear that Φ satisfies conditions (i) and (ii) of Lemma 3.10. To verify the condition (iii) of Lemma 3.10, define for $\psi \in A$, $\varphi \in B$ and $t > 0$:

$$E_t(\psi) := \{x \in X; \psi(x) \leq t\},$$

$$\Phi_t(\psi, \varphi) := \int_{E_t(\psi)} \psi \varphi dm.$$

Then for $\psi \in A$ and $t > 0$, the map $\Phi_t(\psi, \cdot): B \rightarrow \mathbb{R}_+$ is weakly continuous, and for $\varphi \in B$, we have

$$\Phi(\psi, \varphi) = \sup_{t > 0} \Phi_t(\psi, \varphi).$$

This shows that $\Phi(\psi, \cdot)$ is weakly lower-semicontinuous on B . Let $M := \sup_{\psi \in A} \psi^*(\frac{\varepsilon}{2})$, and assume that $M < \infty$ (if $M = \infty$, there is nothing to prove). Then for every $\psi \in A$,

$$\begin{aligned} 1 - \int_X 1_{E_M(\psi)} dm &= m(\{x \in X; |\psi(x)| > M\}) \\ &\leq m(\{x \in X; |\psi(x)| > \psi^*(\frac{\varepsilon}{2})\}) \leq \frac{\varepsilon}{2}. \end{aligned}$$

Here we used Lemma 3.8. This shows that $1_{E_M(\psi)} \in B$ for every $\psi \in A$. Therefore by Lemma 3.10,

$$\begin{aligned} \min_{\varphi \in B} \sup_{\psi \in A} \int_X \psi \varphi dm &= \sup_{\psi \in A} \min_{\varphi \in B} \int_X \psi \varphi dm \\ (12) \qquad \qquad \qquad &\leq \sup_{\psi \in A} \int_{E_M(\psi)} \psi dm \leq M. \end{aligned}$$

Therefore the minimum of the left hand side will be attained at some $\varphi \in B$. So the set B_0 of all such minimizers $\varphi \in B$ is nonempty. We observe that B_0 is a globally Γ -invariant weakly closed, convex subset of B . Indeed, let $0 < \lambda < 1$ and $\varphi_1, \varphi_2 \in B_0$ be minimizers and let $\varphi = \lambda\varphi_1 + (1 - \lambda)\varphi_2$. Then for $\psi \in A$,

$$\begin{aligned} \sup_{\psi \in A} \int_X \psi \varphi dm &\leq \lambda \sup_{\psi \in A} \int_X \psi \varphi_1 dm + (1 - \lambda) \sup_{\psi \in A} \int_X \psi \varphi_2 dm \\ &= \min_{\varphi' \in B} \sup_{\psi \in A} \int_X \psi \varphi' dm. \end{aligned}$$

This shows that φ is also a minimizer, hence in B_0 . Next, let $(\varphi_n)_{n=1}^\infty$ be a sequence of elements in B_0 converging weakly to $\varphi \in L^2(X, m)$. Then $\varphi \in B$, and Fatou's Lemma shows

$$\begin{aligned} \sup_{\psi \in A} \int_X \psi \varphi dm &\leq \sup_{\psi \in A} \liminf_{k \rightarrow \infty} \int_X \underbrace{\psi 1_{\{x; \psi(x) \leq k\}}}_{\in L^2(X, m)} \varphi dm \\ &= \sup_{\psi \in A} \liminf_{k \rightarrow \infty} \lim_{n \rightarrow \infty} \int_X \psi 1_{\{x; \psi(x) \leq k\}} \varphi_n dm \\ &\leq \sup_{\psi \in A} \liminf_{k \rightarrow \infty} \limsup_{n \rightarrow \infty} \int_X \psi \varphi_n dm \\ &\leq \sup_{\psi \in A} \limsup_{n \rightarrow \infty} \sup_{\psi' \in A} \int_X \psi' \varphi_n dm \\ &\stackrel{\varphi_n \in B_0}{=} \limsup_{n \rightarrow \infty} \min_{\varphi' \in B} \sup_{\psi' \in A} \int_X \psi' \varphi' dm \\ &= \min_{\varphi' \in B} \sup_{\psi' \in A} \int_X \psi' \varphi' dm. \end{aligned}$$

This shows that φ is a minimizer too, so $\varphi \in B_0$. Finally, let $\varphi \in B_0$ and $s \in \Gamma$. Then since A is globally Γ -invariant and m is Γ -invariant, we have (recall that the Γ action on X is given

so that $\beta_s(\varphi) = \varphi(s^{-1} \cdot)$

$$\begin{aligned} \sup_{\psi \in A} \int_X \psi \beta_s(\varphi) \, dm &= \sup_{\psi \in A} \int_X \beta_{s^{-1}}(\psi) \varphi \, dm(g \cdot) \\ &= \sup_{\psi \in \beta_{s^{-1}}(A)} \int_X \psi \varphi \, dm \\ &= \min_{\varphi' \in B} \sup_{\psi \in A} \int_X \psi \varphi' \, dm. \end{aligned}$$

This shows that $\beta_s(\varphi) \in B_0$. Therefore B_0 is a Γ -invariant, non-empty, weakly compact, and convex subset of $L^2(X, m)$. Let φ_0 be an element in B_0 of minimum L^2 -norm. By the uniform convexity, such an element is unique. Then since $\|\beta_s(\varphi_0)\|_{L^2(X)} = \|\varphi_0\|_{L^2(X)}$ ($s \in \Gamma$), we have $\beta_s(\varphi_0) = \varphi_0$ ($s \in \Gamma$) by the uniqueness of φ_0 . Then define

$$S(\varepsilon) := \{x \in X; \varphi_0(x) \geq \tfrac{1}{2}\}.$$

Then $S(\varepsilon)$ is a Γ -invariant measurable subset of X . Moreover,

$$\begin{aligned} m(X \setminus S(\varepsilon)) &= m(\{x \in X; \varphi_0(x) < \tfrac{1}{2}\}) \\ &\leq 2 \int_X (1 - \varphi_0(x)) \, dm \leq \varepsilon, \\ \sup_{\psi \in A} \int_{S(\varepsilon)} \psi \, dm &\leq 2 \sup_{\psi \in A} \int_X \psi \varphi_0 \, dm \leq 2M. \end{aligned}$$

Here in the last line we used (12). □

In the sequel, we will mainly follow the arguments in [GR85, Theorem VI.3.3]. Since this source is hardly accessible, we decided to repeat some of the arguments (and present them in a streamlined form) in order to provide a complete proof. We set $Z := \{|(T\xi)(x)|^2; \xi \in H, \|\xi\| \leq 1\} \subset L^0(X, m)_+$ and note that

$$(13) \quad A := \left\{ \sum_{j=1}^n |(T\xi_j)|^2; n \in \mathbb{N}, \xi_1, \dots, \xi_n \in H, \sum_{j=1}^n \|\xi_j\|^2 \leq 1 \right\} \subset L^0(X, m)_+$$

is the convex hull of Z . In order to apply Proposition 3.9 to (a subset of) A , we need to show that the estimates from Lemma 3.7 extend to the convex hull of Z . This is the content of the following proposition.

Proposition 3.11. *There exists a function $(0, \infty) \ni \lambda \mapsto C(\lambda) \in [0, 1]$ satisfying*

$$\lim_{\lambda \rightarrow \infty} C(\lambda) = 0,$$

such that for every finite family $\xi_1, \dots, \xi_n \in H$, one has

$$\sum_{j=1}^n \|\xi_j\|^2 \leq 1 \Rightarrow m \left(\left\{ x \in X; \left(\sum_{j=1}^n |(T\xi_j)(x)|^2 \right)^{\frac{1}{2}} \geq \lambda \right\} \right) \leq C(\lambda).$$

This is Condition (d) in [GR85, Theorem VI.3.3], and the proof can be found in [GR85, Theorem VI.3.6]. We give an alternative and direct proof, following a strategy that we learned from G. Lowther [Low12] on MathOverflow. First, we need a classical Khintchine type inequality in L^0 -space. We fix a *Rademacher sequence* $(\epsilon_n)_{n=1}^\infty$ on a probability space

$(\Omega, \mathbb{P})$. That is, $(\epsilon_n)_{n=1}^\infty$ is a sequence of independent, identically distributed random variables on $(\Omega, \mathbb{P})$ with the distribution $\mathbb{P}(\epsilon_n = 1) = \frac{1}{2} = \mathbb{P}(\epsilon_n = -1)$ ($n \in \mathbb{N}$). The symbol $\mathbb{E}$ denotes the expectation value with respect to $\mathbb{P}$.

Lemma 3.12. *For all $n \in \mathbb{N}$ and for all $x = (x_j)_{j=1}^n \in \mathbb{R}^n$, we have*

$$\mathbb{P} \left(\left(\sum_{j=1}^n \epsilon_j x_j \right)^2 \geq \frac{1}{2} \cdot \sum_{j=1}^n x_j^2 \right) \geq \frac{1}{12}.$$

Proof. Without loss of generality, we may assume that $\sum_{j=1}^n x_j^2 = 1$. We consider the random variable $f(\epsilon_1, \dots, \epsilon_n) = \left(\sum_{j=1}^n \epsilon_j x_j \right)^2$. Then, using independence, we compute $\mathbb{E}(f(\epsilon_1, \dots, \epsilon_n)) = \sum_{j=1}^n x_j^2 = 1$ and

$$\mathbb{E}(f(\epsilon_1, \dots, \epsilon_n)^2) = \sum_{j=1}^n x_j^4 + 3 \sum_{i \neq j} x_i^2 x_j^2 = 3 \left(\sum_{j=1}^n x_j^2 \right)^2 - 2 \sum_{j=1}^n x_j^4 \leq 3.$$

Using the Paley-Zygmund inequality, which asserts that if $Y \geq 0$ is a random variable with $0 \neq \mathbb{E}(Y^2) < \infty$, then

$$\mathbb{P}(Y > \theta \cdot \mathbb{E}(Y)) \geq (1 - \theta)^2 \cdot \frac{\mathbb{E}(Y)^2}{\mathbb{E}(Y^2)} \quad \text{for } 0 \leq \theta \leq 1,$$

we obtain

$$\mathbb{P} \left(f(\epsilon_1, \dots, \epsilon_n) \geq \frac{1}{2} \cdot \mathbb{E}(f(\epsilon_1, \dots, \epsilon_n)) \right) \geq \frac{1}{4} \cdot \frac{\mathbb{E}(f(\epsilon_1, \dots, \epsilon_n))^2}{\mathbb{E}(f(\epsilon_1, \dots, \epsilon_n)^2)} \geq \frac{1}{12}.$$

This proves the claim. $\square$

Proof of Proposition 3.11. Fix $\xi_1, \dots, \xi_n \in H$ with $\sum_{j=1}^n \|\xi_j\|^2 \leq 1$. By Lemma 3.7, there exists a non-increasing function $C': (0, \infty) \rightarrow [0, 1]$, such that $\lim_{\lambda \rightarrow \infty} C'(\lambda) = 0$ satisfying

$$m(\{x \in X; |(T\xi)(x)| \geq \lambda\}) \leq C'(\lambda)$$

for all $\xi \in H$ with $\|\xi\| \leq 1$.

For any constant $\alpha > 0$, we compute

$$\begin{aligned} & \int_X \mathbb{P} \left(\left(\sum_{j=1}^n \epsilon_j T(\xi_j)(x) \right)^2 \geq \alpha \right) dm(x) \\ & \geq \int_{\{x \in X; \frac{1}{2} \cdot \sum_{j=1}^n T(\xi_j)(x)^2 \geq \alpha\}} \mathbb{P} \left(\left(\sum_{j=1}^n \epsilon_j T(\xi_j)(x) \right)^2 \geq \frac{1}{2} \cdot \sum_{j=1}^n T(\xi_j)(x)^2 \right) dm(x) \\ (14) \quad & \geq \frac{1}{12} \cdot m \left(\left\{ x \in X; \frac{1}{2} \cdot \sum_{j=1}^n (T\xi_j(x))^2 \geq \alpha \right\} \right) =: \frac{p(\alpha)}{12}. \end{aligned}$$

Here in the last inequality we used Lemma 3.12. For each $\omega \in \Omega$, define

$$A(\omega) := \left\{ x \in X; \left(\sum_{j=1}^n \epsilon_j(\omega) (T\xi_j)(x) \right)^2 \geq \alpha \right\} \subset X,$$

and set $B := \{\omega \in \Omega; m(A(\omega)) \geq \frac{p(\alpha)}{24}\}$. If $\mathbb{P}(B) < \frac{p(\alpha)}{24}$, then by Fubini Theorem,

$$\begin{aligned} \int_X \mathbb{P} \left(\left(\sum_{j=1}^n \epsilon_j(T\xi_j)(x) \right)^2 \geq \alpha \right) dm(x) &\leq \mathbb{P}(B) + \int_X \mathbb{P}(\omega \in B^c; x \in A(\omega)) dm(x) \\ &< \frac{p(\alpha)}{24} + \int_{B^c} \int_{A(\omega)} dm(x) d\mathbb{P}(\omega) \\ &< \frac{p(\alpha)}{12}, \end{aligned}$$

which contradicts (14). Therefore $\mathbb{P}(B) \geq \frac{p(\alpha)}{24}$ holds. Since H is a Hilbert space, we have the equality

$$\mathbb{E} \left(\left\| \sum_{j=1}^n \epsilon_j \xi_j \right\|^2 \right) = \sum_{j=1}^n \|\xi_j\|^2 \leq 1.$$

Therefore by a similar argument, we can show that the set $C := \{\omega \in \Omega; \|\sum_{j=1}^n \epsilon_j(\omega) \xi_j\|^2 \leq \frac{48}{p(\alpha)}\}$ satisfies $\mathbb{P}(C) \geq 1 - \frac{p(\alpha)}{48}$. In particular, $\mathbb{P}(B \cap C) > 0$ holds (if $p(\alpha) = 0$, then $C = \Omega$, in which case the conclusion is still true). Let $\omega \in B \cap C \neq \emptyset$ and set $\xi = \sum_{j=1}^n \epsilon_j(\omega) \xi_j$. Then $\|\xi\|^2 \leq \frac{48}{p(\alpha)}$, so that

$$\begin{aligned} p(\alpha) = m \left(\left\{ x \in X; \sum_{j=1}^n (T\xi_j(x))^2 \geq 2\alpha \right\} \right) &\leq 24 \cdot m(\{x \in X; (T\xi(x))^2 \geq \alpha\}) \\ &\leq 24 \cdot C' \left(\left(\frac{\alpha \cdot p(\alpha)}{48} \right)^{\frac{1}{2}} \right) \end{aligned}$$

We apply this now with $\lambda^2 = 2\alpha$. If $p(\alpha) = p(\frac{\lambda^2}{2}) > \lambda^{-1}$, then $\alpha p(\alpha) > \frac{\lambda}{2}$ and since $C'(\cdot)$ is non-increasing, we have

$$p(\alpha) = m \left(\left\{ x \in X; \sum_{j=1}^n (T\xi_j(x))^2 \geq \lambda^2 \right\} \right) \leq 24C' \left(\left(\frac{\lambda}{96} \right)^{\frac{1}{2}} \right)$$

We can now set $C(\lambda) := \max \left\{ \lambda^{-1}, 24C' \left(\left(\frac{\lambda}{96} \right)^{\frac{1}{2}} \right) \right\}$. This finishes the proof. $\square$

We are now ready to prove the existence of an equivariant Maurey–Nikishin factorization.

Proof of Theorem 3.5. This part of the proof is exactly the same as Maurey’s argument given as in [GR85, Theorem 3.3]. We include the proof for the reader’s convenience. Define a subset A' of A (see (13)) by

$$A' := \left\{ \sum_{j=1}^n |(T\xi_j)|^2; n \in \mathbb{N}, \xi_1, \dots, \xi_n \in H, \sum_{j=1}^n \|\xi_j\|_\pi^2 \leq 1 \right\} \subset L^0(X, m)_+,$$

where $\|\xi\|_\pi := \sup_{s \in \Gamma} \|\pi(s)\xi\|$ ($\xi \in H$). Then since $\|\pi(s)\xi\|_\pi = \|\xi\|_\pi$ ($s \in \Gamma, \xi \in H$), by (6), the set A' is globally Γ -invariant and convex. Since T is continuous and using Proposition 3.11, a very similar argument as in the proof of Lemma 3.7 (ii) $\Rightarrow$ (iii) implies the following:

for every $t > 0$, there exists $K(t) > 0$ such that for any finite sequence $\xi_1, \dots, \xi_n \in H$, one has (using $(\psi^2)^\star = (\psi^\star)^2$)

$$\left(\sum_{j=1}^n (T\xi_j)^2 \right)^\star(t) \leq K(t)^2 \sum_{j=1}^n \|\xi_j\|^2.$$

By Proposition 3.9, for every $\varepsilon > 0$, there exists a Γ -invariant measurable subset $S(\varepsilon) \subset X$ with $m(X \setminus S(\varepsilon)) < \varepsilon$ satisfying (11) with A replaced by A' . In particular, this shows by (11) that for every $\varphi \in A'$,

$$\int_{S(\varepsilon)} \varphi(x) dm(x) \leq 2K \left(\frac{\varepsilon}{2} \right)^2$$

holds. Therefore for every $\xi \in H$, $\|\xi\|_\pi^2 \leq \|\pi\|^2 \|\xi\|^2$ (where $\|\pi\| := \sup_{s \in \Gamma} \|\pi(s)\|$), so that we have

$$\int_{S(\varepsilon)} [T\xi](x)^2 dm(x) \leq 2K \left(\frac{\varepsilon}{2} \right)^2 \|\pi\|^2 \|\xi\|^2.$$

Set $C_\varepsilon := 2K \left(\frac{\varepsilon}{2} \right)^2 \|\pi\|^2 > 0$. Then let

$$\varphi(x) := \sum_{j=1}^{\infty} k_j 1_{S(1/j)}(x), \quad x \in X,$$

where $k_j > 0$ and $\sum_{j=1}^{\infty} k_j C_{1/j} = 1$. Since C_ε is non-increasing in ε , $k_j \xrightarrow{j \rightarrow \infty} 0$, which implies that $\varphi \in L^\infty(X, m)$. Also, φ is m -a.e. positive and Γ -invariant. Moreover we have:

$$\begin{aligned} \int_X \varphi(x) [T\xi](x)^2 dm(x) &= \sum_{j=1}^{\infty} k_j \int_{S_{1/j}} [T\xi](x)^2 dm(x) \\ &\leq \sum_{j=1}^{\infty} k_j C_{1/j} \|\xi\|^2 = \|\xi\|^2. \end{aligned}$$

This finishes the proof. $\square$

Recall from the non-commutative integration theory that a self-adjoint operator $A \in \widetilde{M}$ is square-integrable, if

$$\|A\|_2^2 := \int_{\mathbb{R}} \lambda^2 d(\tau(E_A(\lambda))) < \infty.$$

Here, $A = \int_{\mathbb{R}} \lambda dE_A(\lambda)$ is the spectral decomposition of A . Then $T \in \widetilde{M}$ is called square-integrable if $|T| = (T^*T)^{\frac{1}{2}}$ is square-integrable. The set of all square-integrable operators can naturally be identified with $L^2(M)$ (see [Ne74] for details). We need the following simple but useful lemma.

Lemma 3.13. *Let $S: H \rightarrow \widetilde{M}$ be an $\mathbb{R}$ -linear homeomorphism onto its range. If $S(\xi) \in L^2(M)$ for every $\xi \in H$, then regarded as $S: H \rightarrow L^2(M)$, S is an $\mathbb{R}$ -linear homeomorphism onto its range, where $L^2(M)$ is equipped with the L^2 -topology (instead of the τ -measure topology).*

Proof. We first show that in $L^2(M)$, the L^2 -topology is stronger than the τ -measure topology. Let $A_n \in L^2(M)$ ($n \in \mathbb{N}$) be such that $\|A_n\|_2 \xrightarrow{n \rightarrow \infty} 0$. Then

$$\begin{aligned} \|(A_n - i)^{-1} - (0 - i)^{-1}\|_2^2 &= \int_{\mathbb{R}} \left| \frac{1}{\lambda - i} - \frac{1}{0 - i} \right|^2 d\tau(E_{A_n}(\lambda)) \\ &= \int_{\mathbb{R}} \frac{\lambda^2}{\lambda^2 + 1} d\tau(E_{A_n}(\lambda)) \\ &\leq \int_{\mathbb{R}} \lambda^2 d\tau(E_{A_n}(\lambda)) \\ &= \|A_n\|_2^2 \xrightarrow{n \rightarrow \infty} 0. \end{aligned}$$

This shows the claim. Note also that this proves that $S^{-1}: S(H) \ni S(\xi) \rightarrow \xi \in H$ is continuous:

$$\begin{aligned} \|S(\xi_n)\|_2 \xrightarrow{n \rightarrow \infty} 0 &\Rightarrow S(\xi_n) \xrightarrow{n \rightarrow \infty} 0 \text{ SRT} \\ &\Leftrightarrow \|\xi_n\| \xrightarrow{n \rightarrow \infty} 0. \end{aligned}$$

Next, we show that $S: H \rightarrow L^2(M)$ is L^2 -continuous. Since $H, L^2(M)$ are Banach spaces, it suffices to show that S is closed, by the Closed Graph Theorem. So assume that $\xi_n \in H$ converges to 0, and $S(\xi_n)$ converges in L^2 to some $A \in L^2(M)$. Then $\|S(\xi_n) - A\|_2 \xrightarrow{n \rightarrow \infty} 0$, whence by the first part, we also have $\overline{S(\xi_n) - A} \xrightarrow{n \rightarrow \infty} 0$ in SRT. Since SRT is a vector space topology in $\widetilde{M}$ (because M is finite), we obtain $S(\xi_n) \xrightarrow{n \rightarrow \infty} A$ in SRT. However, S is continuous in SRT, whence $S(\xi_n) \xrightarrow{n \rightarrow \infty} 0$ in SRT. Since SRT is Hausdorff, $A = 0$. Therefore S is closed, whence L^2 -continuous. $\square$

Remark 3.14. In the above lemma, we did not claim that in $L^2(M)$, SRT and the L^2 -topology are the same. In fact the two topologies are different (but they agree on the image of S , i.e., S defines a strong embedding in the sense of [AK06, Definition 6.4.4]). To see this in the commutative case, let (Ω, μ) be a diffuse probability space, and choose a sequence $(B_n)_{n=1}^\infty$ of measurable subsets of Ω such that $0 < \mu(B_n) \xrightarrow{n \rightarrow \infty} 0$. Define $f_n \in L^2(\Omega, \mu)$ by

$$f_n(x) = \frac{1}{\mu(B_n)^{\frac{1}{2}}} 1_{B_n}(x), \quad x \in \Omega.$$

Then $\|f_n\|_2^2 = 1$ for $n \in \mathbb{N}$. On the other hand,

$$\begin{aligned} \|(f_n - i)^{-1} - (0 - i)^{-1}\|_{L^2(\Omega, \mu)}^2 &= \int_{B_n} \frac{1}{1 + \mu(B_n)} d\mu(x) \\ &\leq \int_{B_n} 1 d\mu(x) = \mu(B_n) \xrightarrow{n \rightarrow \infty} 0 \end{aligned}$$

This shows that $f_n \xrightarrow{n \rightarrow \infty} 0$ (SRT) in $L^0(\Omega, \mu)$, but $\|f_n\|_2 = 1 \not\rightarrow 0$.

Proof of Theorem 3.1. By Theorem 3.5, there exists an m -a.e. positive $\varphi \in L^0(X, m)$ such that (8) holds. We define an $\mathbb{R}$ -bilinear form $\langle \cdot, \cdot \rangle'_\pi: H \times H \rightarrow \mathbb{R}$ by

$$\langle \xi, \eta \rangle'_\pi := \int_X \varphi(x) [T\xi](x) [T\eta](x) dm(x), \quad \xi, \eta \in H.$$

If $\langle \xi, \xi \rangle'_\pi = 0$, then $\varphi |T\xi|^2 = 0$ a.e., and since $\varphi > 0$ a.e., $T\xi = 0$ a.e. But since T is injective, we get $\xi = 0$. Thus this is an inner product. Let $s \in \Gamma$. Then for $\xi, \eta \in H$, the Γ -invariance of φ and m show that

$$\begin{aligned} \langle \pi(s)\xi, \pi(s)\eta \rangle'_\pi &= \int_X \varphi(x) [T\xi](s^{-1}x) [T\eta](s^{-1}x) \, dm(x) \\ &= \int_X \varphi(sx) [T\xi](x) [T\eta](x) \, dm(sx) \\ &= \int_X \varphi(x) [T\xi](x) [T\eta](x) \, dm(x). \end{aligned}$$

Therefore $\langle \cdot, \cdot \rangle'_\pi$ is Γ -invariant. We show that this inner product generates the same topology on H as the original norm. Define $S: H \rightarrow L^2(X, m) \subset L^2(M)$ by $S(\xi) := \varphi^{1/2} T(\xi) = M_{\varphi^{1/2}} T(\xi)$. Note that the multiplication operator $M_{\varphi^{1/2}}$ is a (possibly unbounded) operator affiliated with M . Therefore, the function

$$M_{\varphi^{1/2}}: \widetilde{M} \ni X \mapsto \varphi^{1/2} X \in \widetilde{M}$$

is continuous with continuous inverse $M_{\varphi^{-1/2}}$, so it is a homeomorphism. Therefore we obtain that $S: H \rightarrow \widetilde{M}$ is also an $\mathbb{R}$ -linear homeomorphism onto its range. Moreover, we have the inclusion $S(H) \subset L^2(X, m) \subset L^2(M)$. Thus by Lemma 3.13, $S: H \rightarrow L^2(M)$ is a homeomorphism onto its range where $L^2(M)$ is equipped with the L^2 -topology. Let $(\xi_n)_{n=1}^\infty$ be a sequence in H . Then

$$\|\xi_n\| \xrightarrow{n \rightarrow \infty} 0 \Leftrightarrow \|S(\xi_n)\|_{L^2(M)} = \|\xi\|'_\pi \xrightarrow{n \rightarrow \infty} 0.$$

This shows that $\|\cdot\|'_\pi$ generates the same topology as $\|\cdot\|$. Finally, we complexify the inner product. Define a new sesqui-linear form $\langle \cdot, \cdot \rangle_\pi: H \times H \rightarrow \mathbb{C}$ by

$$(15) \quad \langle \xi, \eta \rangle_\pi := \frac{1}{4} \{ \langle \xi, \eta \rangle'_\pi + \langle i\xi, i\eta \rangle'_\pi + i\langle \xi, i\eta \rangle'_\pi - i\langle i\xi, \eta \rangle'_\pi \}, \quad \xi, \eta \in H.$$

Since $\langle \cdot, \cdot \rangle'_\pi$ is $\mathbb{R}$ -bilinear, so is $\langle \cdot, \cdot \rangle_\pi$. But if $\xi, \eta \in H$, then

$$\begin{aligned} \langle i\xi, \eta \rangle_\pi &= \frac{1}{4} \{ \langle i\xi, \eta \rangle'_\pi + \langle -\xi, i\eta \rangle'_\pi + i\langle i\xi, i\eta \rangle'_\pi + i\langle \xi, \eta \rangle'_\pi \} \\ &= \frac{1}{4} \{ i\langle \xi, \eta \rangle'_\pi + i\langle i\xi, i\eta \rangle'_\pi - \langle \xi, i\eta \rangle'_\pi + \langle i\xi, \eta \rangle'_\pi \} \\ &= i\langle \xi, \eta \rangle_\pi, \\ \langle \xi, i\eta \rangle'_\pi &= \frac{1}{4} \{ \langle \xi, i\eta \rangle'_\pi - \langle i\xi, \eta \rangle'_\pi - i\langle \xi, \eta \rangle'_\pi - i\langle i\xi, i\eta \rangle'_\pi \} \\ &= \frac{1}{4} \{ -i\langle \xi, \eta \rangle'_\pi - i\langle i\xi, i\eta \rangle'_\pi + \langle \xi, i\eta \rangle'_\pi - \langle i\xi, \eta \rangle'_\pi \} \\ &= -i\langle \xi, \eta \rangle_\pi. \end{aligned}$$

Thus $\langle \cdot, \cdot \rangle_\pi$ is sesqui-linear. Note that

$$\|\xi\|_\pi^2 = \frac{1}{4} (\|\xi\|'_\pi)^2 + \frac{1}{4} (\|i\xi\|'_\pi)^2,$$

so that $\langle \cdot, \cdot \rangle_\pi$ also generates the same Hilbert space topology. Since $\pi(s)$ ($s \in \Gamma$) is $\mathbb{C}$ -linear and preserves $\langle \cdot, \cdot \rangle'_\pi$, it is also clear that $\pi(s)$ preserves $\langle \cdot, \cdot \rangle_\pi$ as well. By Lemma 3.2, this shows that π is unitarizable. $\square$

Note that in our construction $G = H \rtimes_{\pi} \Gamma$, it is essential that Γ is discrete. It is therefore of interest to consider the following question:

Question 3.15. Is there a connected, unitarily representable SIN Polish group which is not of finite type?

Note also that in the proof of Theorem 3.1, we have shown the following result, which is of independent interest:

Corollary 3.16. *Let H be a separable real Hilbert space, and let $f \in \mathcal{P}(H)$. Then there exist*

- (i) *a compact metrizable space X and a Borel probability measure m on X ,*
- (ii) *a bounded operator $T: H \rightarrow L^2(X, m; \mathbb{R})$, and*
- (iii) *a positive measurable function $\varphi \in L^\infty(X, m)$,*

such that

$$f(\xi) = \int_X e^{i\varphi(x)[T\xi](x)} dm(x), \quad \xi \in H.$$

Moreover:

- (iv) *If f generates a neighborhood basis of 0, then T can be chosen to be a real-linear homeomorphism onto its range.*
- (v) *We may arrange T and φ so that if $\mathcal{G} = \{u \in O(H); f(u\xi) = f(\xi) \ (\xi \in H)\}$ is the subgroup of the orthogonal group of H fixing f pointwise, then there exists a continuous homomorphism β from $\mathcal{G}$ to the group $\text{Aut}(X, m)$ of all m -preserving automorphisms of X with the weak topology, such that*

$$[T(u\xi)](x) = \beta(u)[T(\xi)](x), \quad \varphi(\beta(u)x) = \varphi(x), \quad u \in \mathcal{G}, \ \xi \in H, \ m\text{-a.e. } x \in X.$$

One may regard Corollary 3.16 as an equivariant version of the classical result (cf. [AMM85, Lemma 4.2]) that any $f \in \mathcal{P}(H)$ is of the form

$$f(t\xi) = \int_X e^{it[U\xi](x)} dm(x), \quad \xi \in H, \ t \in \mathbb{R}$$

for a probability space (X, m) and a continuous linear map $U: H \rightarrow L^0(X, m)$.

Remark 3.17. It is clear from the proof that it applies whenever H is a Banach space of Rademacher type 2 (see [AK06, §6] for definitions). Thus we have shown as a corollary that a Banach space B is isomorphic (as a Banach space) to a Hilbert space if and only if a positive definite function generates the topology of B and is of type 2. Moreover, the isomorphism can be made equivariant with respect to a group of symmetries that preserve the positive definite function. One may compare this result with Kwapien's Theorem [Kw72] that B is isomorphic to a Hilbert space if and only if B is of type 2 and cotype 2. Note that the Kwapien Theorem does not produce an equivariant isomorphism but requires less in the sense that any Banach space isomorphic to a subspace of $L^0(X, m)$ is of cotype 2, but not all cotype 2 Banach spaces can be isomorphic to a subspace of $L^0(X, m)$ (see the remark after [BL00, Corollary 8.17] and [Pi78]).

APPENDIX

The following two results are well-known, but we include the proofs for the reader's convenience.

Proposition 3.18. *Let A be a commutative von Neumann algebra with separable predual, and let τ be a normal faithful tracial state on A . Let G be a Polish group, and let $\beta: G \rightarrow \text{Aut}(A)$ be a continuous τ -preserving action. Then there exists a separable compact metrizable space X , a Borel probability measure m on X , an m -preserving continuous action $G \times X \ni (g, x) \mapsto gx \in X$ and a $*$ -isomorphism $\Phi: A \rightarrow L^\infty(X, m)$ such that*

$$(16) \quad \tau(a) = \int_X \Phi(a)(x) dm(x), \quad a \in A,$$

$$(17) \quad \Phi(\beta_g(a))(x) = \Phi(a)(g^{-1}x), \quad g \in G, \mu\text{-a.e. } a \in A.$$

Moreover, the G -action is continuous as a map $G \rightarrow \text{Aut}(X, m)$, where the group $\text{Aut}(X, m)$ of all m -preserving automorphism of X is equipped with the weak topology.

Lemma 3.19. *Let M, N be von Neumann algebras, and let ψ, φ be normal faithful states on M, N , respectively. Let M_0 (resp. N_0) be a $*$ -strongly dense subalgebra of M (resp. N). If $\alpha: M_0 \rightarrow N_0$ is a $*$ -isomorphism such that $\varphi(\alpha(x)) = \psi(x)$ for every $x \in M_0$, then α can be extended to a $*$ -isomorphism from M onto N such that $\varphi(\alpha(x)) = \psi(x)$ ($x \in M$).*

Proof. By the GNS construction, we may assume that M acts on $L^2(M, \psi)$ and N acts on $L^2(N, \varphi)$ so that $\psi = \langle \cdot, \xi_\psi \rangle$, $\varphi = \langle \cdot, \xi_\varphi \rangle$ for unit vectors $\xi_\psi \in L^2(M, \psi)$ and $\xi_\varphi \in L^2(N, \varphi)$. Define $U_0: M_0\xi_\psi \rightarrow N_0\xi_\varphi$ by

$$U_0x\xi_\psi := \alpha(x)\xi_\varphi, \quad x \in M_0.$$

Since ψ is faithful, U_0 is well-defined. Also, since M_0 is $*$ -strongly dense in M and ξ_ψ is cyclic for M , $M_0\xi_\psi$ is dense in $L^2(M, \psi)$. Similarly, $N_0\xi_\varphi$ is dense in $L^2(N, \varphi)$. By $\varphi \circ \alpha = \psi$ on M_0 , we have

$$\|U_0x\xi_\psi\|^2 = \varphi(\alpha(x^*x)) = \psi(x^*x) = \|x\xi_\psi\|^2, \quad x \in M_0.$$

Therefore, U_0 extends to an isometry U from $L^2(M, \psi)$ into $L^2(N, \varphi)$ and the range of U contains a dense subspace $N_0\xi_\varphi$. Therefore U is onto. If $x \in M_0$, then for each $y \in N_0$,

$$UxU^*y\xi_\varphi = Ux\alpha^{-1}(y)\xi_\psi = \alpha(x)y\xi_\varphi,$$

so that by the density of $N_0\xi_\varphi$, we have

$$UxU^* = \alpha(x), \quad x \in M_0.$$

Let now $x \in M$. Then there exists a net $(x_i)_{i \in I}$ in M_0 converging $*$ -strongly to x . Then $Ux_iU^* = \alpha(x_i)$ ($\in N$) converges $*$ -strongly to UxU^* . Therefore $UxU^* \in N$. Then it is clear that $M \ni x \mapsto UxU^* \in N$ defines a $*$ -isomorphism with inverse $N \ni y \mapsto U^*yU \in M$. The uniqueness of the extension is clear by the $*$ -strong density of M_0 and the fact that isomorphisms between von Neumann algebras are normal. It is now clear that $\varphi \circ \alpha = \psi$ on M . This finishes the proof. $\square$

Proof of Proposition 3.18. Let $A_0 := \{x \in A; \lim_{g \rightarrow 1} \|\alpha_g(x) - x\| = 0\}$, which is a norm-dense C^* -subalgebra of A . Let $\{a_n\}_{n=1}^\infty$ be a $*$ -strongly dense subset of A_0 , and let $C = C^*(\{\beta_g(a_n); g \in G, n \in \mathbb{N}\} \cup \{1\})$. Then since G is separable and $g \mapsto \beta_g(a_n)$ is norm-continuous for all $n \in \mathbb{N}$, C is a norm-separable $*$ -strongly dense C^* -subalgebra of A , and let $X = \text{Spec}(C)$ (the Gelfand spectrum), which is a separable and compact metrizable space, and let $\Phi: C \rightarrow C(X)$ be the Gelfand transform. Since $\tau \circ \Phi^{-1} \in C(X)_+^*$, there exists a Borel probability measure m on X such that

$$\tau(c) = \int_X \Phi(c)(x) dm(x), \quad c \in C.$$

We let $C(X)$ act on $L^2(X, m)$ by multiplication, so $C(X) \subset L^\infty(X, m)$ is a $*$ -strongly dense $*$ -subalgebra of $L^\infty(X, m)$. The map $\tau_m: L^\infty(X, m) \ni f \mapsto \int_X f \, dm \in \mathbb{C}$ is a normal faithful tracial state on $L^\infty(X, m)$. Moreover, $\tau_m(\Phi(c)) = \tau(c)$ for every $c \in C$. Therefore by Lemma 3.19, Φ can be extended to a $*$ -isomorphism $\Phi: C'' = A \rightarrow C(X)'' = L^\infty(X, m)$ such that $\tau_m(\Phi(a)) = \tau(a)$ for every $a \in A$. This shows (16). Now let $g \in G$ and $x \in X$. Let $m_x \in C(X)^*$ be the delta probability measure at $x \in X$. Let β^* be the dual action of G on $C(X)^*$ given by

$$(18) \quad \langle \beta_g^*(\nu), f \rangle := \langle \nu, \Phi \circ \beta_{g^{-1}} \circ \Phi^{-1}(f) \rangle, \quad \nu \in C(X)^*, f \in C(X), g \in G.$$

Since β is an action, β_g^* maps pure states (on $C \cong C(X)$) to pure states, and pure states on $C(X)$ are precisely the delta probability measures, there exists a unique point $g \cdot x \in X$ such that

$$\beta_g^*(m_x) = m_{g \cdot x}.$$

We show that $G \times X \ni (g, x) \mapsto g \cdot x \in X$ is a continuous m -preserving action. It is routine to check that this is indeed an action of G on X . We show that the action is continuous. Since G is a Polish group and X is a Polish space, it suffices to show that the action is separately continuous (see e.g., [Kec96, Theorem 9.14]). Suppose that $(x_n)_{n=1}^\infty \subset X$ converges to $x \in X$ and let $g \in G$. Then by the definition of the Gelfand spectrum, $m_{x_n} \xrightarrow{n \rightarrow \infty} m_x$ (weak*). Since β_g induces an action on C (C is β -invariant by construction), we have

$$\beta_g^*(m_{x_n}) = m_{g \cdot x_n} \xrightarrow{n \rightarrow \infty} \beta_g^*(m_x) = m_{g \cdot x},$$

which implies $g \cdot x_n \xrightarrow{n \rightarrow \infty} g \cdot x$ (again by the definition of the topology on the Gelfand spectrum). Let now $x \in X$ and $(g_n)_{n=1}^\infty \subset G$ be a sequence converging to $g \in G$. Then for each $f \in C(X)$,

$$\begin{aligned} \langle \beta_{g_n}^*(m_x), f \rangle &= \langle m_x, \Phi \circ \beta_{g_n^{-1}} \circ \Phi^{-1}(f) \rangle \xrightarrow{n \rightarrow \infty} \langle m_x, \Phi \circ \beta_{g^{-1}} \circ \Phi^{-1}(f) \rangle \\ &= \langle \beta_g^*(m_x), f \rangle, \end{aligned}$$

whence $g_n \cdot x \xrightarrow{n \rightarrow \infty} g \cdot x$. Therefore the G -action on X is continuous. Next, we show that for each $g \in G$, the map $X \ni x \mapsto g \cdot x \in X$ preserves m . It suffices to show that

$$\int_X f(g^{-1}x) \, dm(x) = \int_X f(x) \, dm(x), \quad f \in C(X).$$

Let $\nu = \sum_{i=1}^n \lambda_i m_{x_i} \in \text{conv}\{m_x; x \in X\} =: \mathcal{D}$. Then

$$\beta_g^*(\nu) = \sum_{i=1}^n \lambda_i \delta_{g \cdot x_i} = \nu(g^{-1} \cdot).$$

By the Krein-Milman Theorem, $\mathcal{D}$ is weak*-dense in the state space of $C(X)$. This shows that

$$(19) \quad \beta_g^*(\nu) = \nu(g^{-1} \cdot), \quad \nu \in C(X)_+^*.$$

In particular, by $\tau \circ \beta_g = \tau$,

$$\begin{aligned}
\int_X f(g^{-1}x) dm(x) &= \int_X f(x) dm(g(x)) \stackrel{(19)}{=} \langle \beta_{g^{-1}}^*(m), f \rangle \\
&= \langle m, \Phi \circ \beta_g \circ \Phi^{-1}(f) \rangle \\
&= \tau_m(\Phi \circ \beta_g \circ \Phi^{-1}(f)) = \tau(\beta_g \circ \Phi^{-1}(f)) \\
&= \tau(\Phi^{-1}(f)) = \tau_m(f) \\
&= \int_X f(x) dm(x).
\end{aligned}$$

Therefore the G -action on X is m -preserving. Finally, we show (17). Since the map

$$\beta'_g: L^\infty(X, m) \ni f \mapsto f(g^{-1} \cdot) \in L^\infty(X, m)$$

defines a τ_m -preserving action and since $C(X)$ is $*$ -strongly dense in $L^\infty(X, m)$, it suffices to show that (17) holds for every $f = \Phi(a) \in C(X)$ ($a \in C$), $x \in X$, and $g \in G$. But

$$\begin{aligned}
f(g^{-1}x) &= \langle \mu_{g^{-1} \cdot x}, f \rangle = \langle \beta_{g^{-1}}^*(\mu_x), f \rangle \\
&= \langle \mu_x, \Phi \circ \beta_g \circ \Phi^{-1}(f) \rangle \\
&= \Phi \circ \beta_g \circ \Phi^{-1}(f)(x).
\end{aligned}$$

Finally, assume that $g_n \xrightarrow{n \rightarrow \infty} g$ in G . Then for every measurable set $A \subset X$, we have

$$\begin{aligned}
m(g_n A \triangle g A) &= \int_X |\Phi \circ \beta_{g_n^{-1}} \circ \Phi^{-1}(1_A)(x) - \Phi \circ \beta_{g^{-1}} \circ \Phi^{-1}(1_A)(x)| dm(x) \\
&\leq \|\beta_{g_n^{-1}} \circ \Phi^{-1}(1_A) - \beta_{g^{-1}} \circ \Phi^{-1}(1_A)\|_2 \xrightarrow{n \rightarrow \infty} 0,
\end{aligned}$$

because the u -topology on $\text{Aut}(A)$ is the same as the topology of pointwise $\|\cdot\|_2$ -convergence. Thus the G -action defines a continuous homomorphism $G \rightarrow \text{Aut}(X, m)$. This finishes the proof. $\square$

ACKNOWLEDGMENTS

We thank Professors Łukasz Grabowski, Magdalena Musat, Yuri Neretin, Izumi Ojima and Mikael Rørdam for helpful discussions on the $\mathcal{U}_{\text{fin}}$ -problem at early stages of the project. The final part of the manuscript was completed while HA was visiting the Mittag-Leffler Institute for the program “Classification of operator algebras: complexity, rigidity, and dynamics”. He thanks the institute and the workshop organizers for the invitation and for their hospitality. This research was supported by JSPS KAKENHI 16K17608 (HA), 6800055 and 26350231 (YM), ERC Starting Grant No. 277728 (AT) and the Danish Council for Independent Research through grant no. 10-082689/FNU. (AT), from which HA was also supported while he was working at University of Copenhagen.

REFERENCES

- [AMM85] I. Aharoni, B. Maurey and B. S. Mityagin, Uniform embeddings of metric spaces and of Banach spaces into Hilbert spaces, *Israel J. Math.* **52** (1985), 251–265.
- [AK06] F. Albiac and N. Kalton, Topics in Banach space theory, Graduate Texts in Mathematics **233**, Springer (2006).
- [AM12-1] H. Ando and Y. Matsuzawa, Lie Group-Lie Algebra Correspondences of Unitary Groups in Finite von Neumann Algebras, *Hokkaido Math. J.* **41** (2012), 31–99.

- [AM12-2] H. Ando and Y. Matsuzawa, On Polish Groups of Finite Type, *Publ. RIMS* vol. **48** (2012), 389–408.
- [Be10] D. Beltita, Lie theoretic significance of the measure topologies associated with a finite trace, *Forum Math.* **22** (2010), 241–253.
- [BL00] Y. Benyamini and J. Lindenstrauss, Geometric nonlinear functional analysis Vol. **1**, American Mathematical Society Colloquium Publications, **48**. American Mathematical Society, Providence, RI (2000).
- [Bo87] M. Bozejko, Uniformly bounded representations of free groups, *J. Reine Angew. Math.* **377** (1987), 170–186.
- [Da50] M. Day, Means for the bounded functions and ergodicity of the bounded representations of semi-groups, *Trans. Amer. Math. Soc.* **69** (1950), 276–291.
- [Di50] J. Dixmier, Les moyennes invariantes dans les semi-groupes et leurs applications, *Acta Sci. Math. Szeged* **12** (1950) 213–227.
- [EM55] L. Ehrenpreis and F. Mautner, Uniformly bounded representations of groups, *Proc. Nat. Acad. Sci. U. S. A.* **41** (1955), 231–233.
- [EM09] I. Epstein and N. Monod, Nonunitarizable representations and random forests, *Int. Math. Res. Not. IMRN* **22** (2009), 4336–4353.
- [Gal09] J. Galindo, On unitary representability of topological groups, *Math. Z.* **263** (2009), 211–220.
- [Gao05] S. Gao, Unitary group actions and Hilbertian Polish metric spaces, *Contemp. Math.* **380**, Amer. Math. Soc., Providence, RI, 2005.
- [GR85] J. García-Cuerva and J. L. Rubio de Francia, Weighted norm inequalities and related topics, North-Holland Mathematics Studies **116** (1985).
- [Kec96] A. S. Kechris, Classical descriptive set theory, Springer (1996).
- [Kw72] Isomorphic characterizations of inner product spaces by orthogonal series with vector valued coefficients, *Studia Math.* **44** (1972), 583–595.
- [Low12] G. Lowther, a comment on Mathoverflow at <http://mathoverflow.net/questions/98410>
- [MZ83-1] A. M. Mantero and A. Zappa, The Poisson transform and representations of a free group, *J. Funct. Anal.* **51** (1983), 372–399.
- [MZ83-2] A. M. Mantero and A. Zappa, Uniformly bounded representations and L^p -convolution operators on a free group, Harmonic analysis (Cortona, 1982), 333–343, Lecture Notes in Math., **992**, Springer (1983).
- [Ma72] B. Maurey, Théorèmes de Nikishin: théorèmes de factorisation pour les applications lineaires a valeurs dans un espace $L^0(\Omega, \mu)$, *Seminaire d'analyse fonctionnelle (Polytechnique)* **10** and **11** (1972–1973), 1–10.
- [Me02] M. Megrelishvili, Reflexively but not unitarily representable topological groups. *Topology Proc.* **25** (2002), 615–625.
- [MO10] N. Monod and N. Ozawa, The Dixmier problem, lamplighters and Burnside groups, *J. Funct. Anal.* **258** no. 1 (2010), 255–259.
- [Ne74] E. Nelson, Notes on non-commutative integration, *J. Funct. Anal.* **15** (1974), 103–116.
- [Ni70] E. M. Nikishin, Resonance theorems and superlinear operators (Russian), *Uspehi Mat. Nauk* **25** (156) (1970), 129–191.
- [NT51] M. Nakamura and Z. Takeda, Group representation and Banach limit, *Tohoku Math. J.* **3** (1951), 132–135.
- [Os09] D. Osin, L^2 -Betti numbers and non-unitarizable groups without free subgroups, *Int. Math. Res. Not. IMRN* **22** (2009), 4220–4231.
- [Oz06] N. Ozawa, An Invitation to the Similarity Problems (after Pisier) (2006), available at <http://www.kurims.kyoto-u.ac.jp/~narutaka/notes/similarity.pdf>
- [Po07] S. Popa, Cocycle and orbit equivalence superrigidity for malleable actions of w -rigid groups, *Invent. Math.* **170** (2007), 243–295.
- [Pi78] G. Pisier, Some results on Banach spaces without local unconditional structure, *Compositio Math.* **37** (1978), 3–19.
- [Pi86] G. Pisier, Factorization of linear operators and geometry of Banach spaces, CBMS Regional Conference Series in Mathematics, vol. **60**, Published for the Conference Board of the Mathematical Sciences, Washington, DC (1986).
- [Pi01] G. Pisier, Similarity problems and completely bounded maps. Second, Lecture Notes in Mathematics, **1618**. Springer-Verlag, (2001).

- [Pi05] G. Pisier, Are unitarizable groups amenable?, in: *Infinite groups: geometric, combinatorial and dynamical aspects*, Progress in Mathematics **248**, Birkhäuser (2005), 323–362.
- [PS86] T. Pitlyk and R. Szwarc, An analytic family of uniformly bounded representations of free groups, *Acta. Math.* **157**, 12 (1986), 287–309.
- [SzN47] B. Sz.-Nagy, On uniformly bounded linear transformations in Hilbert space, *Acta Sci. Math.* (Szeged) **11** (1947), 152–157.
- [Sz88] R. Szwarc, An analytic series of irreducible representations of the free group, *Ann. Inst. Fourier* (Grenoble) **38** (1988), 87–110.
- [VZ74] F.H. Vasilescu and L. Zsido, Uniformly bounded groups in finite W^* -algebras, *Acta Sci. Math.* (Szeged), **36** (1974), 189–192.
- [Ya85] Y. Yamasaki, Measures on infinite dimensional spaces, Series in Pure Mathematics vol. **5**, World Scientific (1985)
- [Ye71] F. J. Yeadon, A new proof of the existence of a trace in a finite von Neumann algebra, *Bull. Amer. Math. Soc.* **77** (1971), 257–260.

HIROSHI ANDO, DEPARTMENT OF MATHEMATICS AND INFORMATICS, CHIBA UNIVERSITY, 1-33 YAYOI-CHO, INAGE, CHIBA, 263-0022, JAPAN

E-mail address: `hiroando@math.s.chiba-u.ac.jp`

YASUMICHI MATSUZAWA, DEPARTMENT OF MATHEMATICS, SHINSHU UNIVERSITY, 6 -RO, NISHINAGANO, NAGANO CITY 380-8544, JAPAN

E-mail address: `myasu@shinshu-u.ac.jp`

ANDREAS THOM, FACHBEREICH MATHEMATIK, INSTITUT FÜR GEOMETRIE, TU DRESDEN, 01062 DRESDEN, GERMANY

E-mail address: `andreas.thom@tu-dresden.de`

ASGER TÖRNQUIST, DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF COPENHAGEN, UNIVERSITETSPARK 5, 2100 COPENHAGEN, DENMARK

E-mail address: `asgert@math.ku.dk`